



Prevalence of Malaria Infection and Molecular Plasmodium Speciation in Subjects Attending Secondary Health Centres in Adamawa State, Nigeria

Angaticha Richard ^{a*}, Nduka Florence Onyemachi ^a
and Abah Austin Edache ^a

^a Department of Animal and Environmental Biology, University of Port Harcourt, P.M.B. 5323 Port Harcourt, Rivers State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117562>

Original Research Article

Received: 03/04/2024

Accepted: 06/06/2024

Published: 11/06/2024

ABSTRACT

Background: Malaria remains a disease of concern as it continues to plunge many into economic difficulty and much suffering, claiming lives especially children under 5 years old. The study investigated prevalence of malaria infection and molecular identification of Plasmodium species in Adamawa state, Nigeria.

Methods: Health facility-based cross-sectional study was conducted from March, 2021 to August, 2021 on subjects attending 15 secondary health facilities distributed across 3 zones in Adamawa State, Nigeria. The prevalence of Plasmodium infection was measured by light microscopic and polymerase chain reaction (RT-PCR) targeting *Plasmodium falciparum* 18S rRNA gene. Infection in relation to location, age, sex and socio-economic status was investigated. The ownership and usage of insecticidal treated mosquito nets (ITNs) were also assessed.

*Corresponding author: E-mail: angaticha@yahoo.com;

Cite as: Richard, Angaticha, Nduka Florence Onyemachi, and Abah Austin Edache. 2024. "Prevalence of Malaria Infection and Molecular Plasmodium Speciation in Subjects Attending Secondary Health Centres in Adamawa State, Nigeria". *South Asian Journal of Parasitology* 7 (3):204-16. <https://journalsajp.com/index.php/SAJP/article/view/185>.

Results: Prevalence by microscopic analysis was 39.08% with total parasite density 1633048/ μ l, while PCR assay amplifying 18S small-subunit ribosomes RNA (SSU rRNA) gene of Plasmodium confirmed only 15.7% of isolate as asymptomatic malaria infections. *P. falciparum* was the only species found in the study area. Infection by geographical locations, age, sex, socio-economic status varied significantly ($P=0.05$). Children under 5 years of age and female subjects recorded highest infection rate by both microscopy and PCR. Shockingly only 52.3% of subjects without ITNs were infected.

Conclusion: The results indicated the need for scale-up interventions to curb the high prevalence of *P. falciparum* infection in the study area with special attention to children and female subjects in the population.

Keywords: Malaria infection; socioeconomic status; molecular identification; Adamawa state.

1. INTRODUCTION

Malaria, though preventable and treatable remained a life-threatening disease caused by protozoan parasite species (*Plasmodium falciparum*, *Plasmodium vivax*, *Plasmodium ovale*, *Plasmodium malariae* and *Plasmodium knowlesi*), transmitted through the bite of an infected female Anopheles mosquito [1]. It is often referred to as disease of the poor due to its socioeconomic relevance in distribution [2,3,4]. Low socioeconomic households suffer high malaria infection due to their inability to prevent and treat malaria, while keeping them struggling to improve their socioeconomic status [4]. The population at risk of contracting malaria and developing severe diseases than others are infants, children under 5 years of age, pregnant women and patients with HIV/AIDS as well as non-immune migrants. Symptoms of severe ill-health from malaria include headache, fever, tiredness and fatigue, convulsion, difficulty in breathing, dark or bloody urine, jaundice and abnormal bleeding [5].

Globally, there has been a rise in malaria cases from 216 million cases (in 91 countries) reported the year 2018 to 247 million cases in 2021 (across 84 malaria endemic countries). Most of the increase was in WHO Africa region and was attributed to the covid-19 era [6, 7]. Among four countries that accounted for half of the world malaria cases Nigeria leads with 27% followed by the Democratic Republic of Congo 12%, Uganda 5% and Mozambique 4% [6]. Hence, it is of no surprise that 97% of Nigerians are at risk of contracting malaria [8].

Plasmodium falciparum malaria has been reported as the leading cause of malaria deaths and burden in sub-Saharan Africa. Falciparum malaria case in Nigeria recently rose from 1.521.566 in 2018, 3.659.170 in 2020 to 3.828.757 in 2021 [7]. WHO in 2019 included

'high political will to reduce malaria' as one of the key elements of high impact approach. Nigeria, in her effort to end malaria by 2030 launched 'End Malaria Council' to assist the Malaria Elimination Program achieves its goal in the country [9].

Preventive measures such as the sleeping under long lasting insecticidal nets (LLINs), insecticide treated mosquito nets (ITNs) and the use of indoor residual spraying (IRS) to control vectors and chemoprevention cannot be overemphasized as effective tools [5]. Across the 21 LGAs in Adamawa State about one million children (aged 3-59 months) benefited from the Seasonal Malaria Chemoprevention (with sulfadoxine-pyremethamine and Amodiaquine) programme with support from WHO and funding from Global Funds through the National Malaria Elimination Programme (NMEP) in 2022 [10]. In 2021, WHO recommended the use of RTS,S/AS01 vaccine for children as it has been found to give protection against malaria infection. About 2 million children have been reached in Kenya, Malawi and Ghana since 2019 with the RTS.S/AS01. In 2023 R21Matrix M was approved as the second vaccine for malaria WHO as it has shown to be effective [11]. Nigeria grants approval for RTS/S vaccine in 2023 which is expected from Oxford [12].

To report recent findings that will pave way for evidence-based decision on effort to achieve goal of ending malaria sufferings in Nigeria, this study investigated the malaria parasitaemia and molecular identification of Plasmodium species in Adamawa State.

2. METHODS

2.1 Study Areas

Adamawa state is located in North Eastern part of Nigeria. It was carved out of the former

Gongola state in 1991 with its headquarters in Yola. It is bordered by Borno and Yobe states in the North, Gombe state in the West, Taraba state in the South and Republic of Cameroun from the East (along Nigerian international border). It has 3 zones (North, Central and South) and 21 local government areas with total population of 3178950 based on the last census conducted in 2006, the projected population in 2022 was 4902100. The minimum and maximum temperatures are 15.2°C to 43°C respectively [13,14]. The annual rainfall ranges from 79mm to 179mm within the period of May to September. Its coordinates are 9°20'N 12°30'E and it has landmass of 36917km² [14,15]. The topography is made up of mountainous land crossed by river valleys of Benue, Gongola and Yadserem. The valleys of Cameroun, Mandara and Adamawa mountains form part of the landscape [16].

The major occupation of the people is farming as reflected in the two vegetation zones; Sudan and Northern Guinea Savannah. The common crops cultivated are groundnuts, cotton, maize, guinea corn, millet, cassava, yam and rice. Communities along the river banks engage in fishing while most of the Fulanis practice nomadic farming [16,17].

2.2 Study Design

2.2.1 Sampling technique

Random sampling technique was used. The study area was further stratified into 3 Zones (North, Central and South) in terms of location. The Northern zone consist of Madagali LGA, Michika LGA, Mubi North LGA, Mubi South LGA, Maiha LGA while Central zone consist of Hong LGA, Gombi LGA, Song LGA, Yola North and




Fig. 1. Map of Nigeria showing study areas
 Source: Department of Geography, University of Port Harcourt

Fufore LGA. Selected from South were Numan LGA, Mayo-Belwa LGA, Demsa LGA, Ganye LGA and Toungo LGA. Total of 15 secondary health facilities were selected; 5 health facilities from each zone. Sample size was determined by Yamane's formula: $n=N/1+N(e)^2$ Where n =sample size, N =population size, e = level of precision. The confidence level at 95% and level of precision used was 0.05. The population size of the 3 zones of Adamawa State was used to determine the total sample size. For example: Adamawa North; population size by 2022 projection is 1051900 and by 2006 census is 682026; applying the Yamane's formula you will have 400. And so, for other zones. Hence $400 \times 3=1200$ [18]. A total of 1200 samples were collected.

2.2.2 Inclusion criteria

The study was delimited to subjects of all ages that presented at the selected secondary health centres irrespective of their social status and those whose parents/guardian signed the consent forms for under children after clear explanation of study objectives both in English and Hausa languages.

2.2.3 Exclusion criteria

Subjects who are on malaria treatment and visitors (e.g. travellers) who report to health facilities and thereafter will leave the study areas were excluded.

2.2.4 Questionnaire administration

The questionnaire was administered through face-to-face interview, however, those that could not express themselves in English language were interviewed in Hausa (Most residents speak the language). Subjects' bio-data and information on socioeconomic status, ownership and usage of ITNs and other variables associated with symptomatic infection were obtained. Each sampling points coordinates was obtained using GPS Application by Ketan computers [19].

2.3 Sample Collection and Storage

For laboratory investigation 5ml of blood was collected from each subject with assistance of the licensed laboratory scientist working in the secondary health facilities, blotted in triplicate onto Whatman 3 paper and allowed to air dry at room temperature. Whatman paper was

individually placed in a sealed plastic bags marked with subject's study numbers and date of collection before being transported to the Nigerian Institute of Medical Research (NIMR) Lagos, Nigeria. The 108 samples for molecular identification were randomly selected across the study areas. For microscopy, films were made at the laboratories of the health facility stained and transported to laboratories in General Hospitals (GH) Mubi and Ganye for investigation.

2.4 Processing of Blood Smears/ Microscopy for Parasite Status and Density

Thick (stained with Field stain A and B) and thin (stained with 3% Giemsa) blood films were prepared and examined by oil immersion microscopy. Thick films are for detection of parasitaemia and thin films for species differentiation. Results obtained was recorded against each questionnaire and compiled in record book. Parasitaemia was calculated using the formular;

$$\text{Parasite count} \times 8000/\text{set range of WBC}=\text{parasite }/\mu\text{l [20].}$$

2.5 Data Analysis

Chi-square test, analysis of variance and descriptive statistics were used for comparison of malaria parasitaemia in the study areas; the package used was statistical package for social science (SPSS) version 2022.

2.6 DNA Extraction

Parasite DNA was extracted from the dried blood spot according to the manufacturer's instructions. The prepared samples were kept at -20°C until use.

2.7 Plasmodium Speciation by Species-Specific Nested PCR

The standard method for molecular identification of *Plasmodium* species carried out relies on ribosomal RNA gene targets in the 18s subunit and requires a nested PCR [21].

Two genus-specific primers rPLU5 and rPLU6 were used for the first cycle amplification. An aliquot of the product thus obtained was used for second amplification cycle in which the individual parasite species was detected

separately using the species-specific primers (Table 1).

A master mix (Taq polymerase enzyme, buffer, magnesium chloride) was prepared for one 12.5µl PCR reactions and reagents were added in order presented in Table 2. Amplification was done following conditions in Table 1. After

amplification of DNA template for Nest 1, Nest 2 reagents were added to PCR tube as provided in Table 2. The mixture was gently mixed and amplification done following condition stated in Table 1. The PCR product (DNA amplified) obtained were subjected to amplicon separation in a 1.5% agarose gel (with ethidium bromide stain) electrophoresis.

Table 1. Details of primer and reaction conditions used for the nested PCR analysis

Primer	Sequence (5'-3')	Cycling condition	Product size
Nest 1	rPLU5 CTTGTTGTTGCCTTAACTTC	1.2kb	
	rPLU6 TTAAAATTGTTGCAGTTAAAACG	94°C-3mins;{95°C-30s;49°C-30s; 72°C-90s}X30; 72°C-5mins	
Nest 2	rFAL1 TTAAACTGGTTTGGGAAAACCAAATATATT	205bp	
	rFAL2 ACACAATGAACTCAATCATGACTACCCGTC	94°C-3mins;{95°C-30s;60°C-30s; 72°C-90s}X30; 72°C-5mins	
Nest 2	rMAL1 ATAACATAGTTGTACGTTAAGAATAACCGC	144bp	
	rMAL2 AAAATTCCCATGCATAAAAAATTATACAAA	94°C-3mins;{95°C-30s;49°C-30s; 72°C-90s}X30; 72°C-5mins	
Nest 2	rVIV1 CGACTTCCAAGCCGAAGCAAAGAAAG	120bp	
	rVIV2 TCCTTACTTCTAGCTTAATCCACATAACTGATAC	94°C-3mins;{95°C-30s;49°C-30s; 72°C-90s}X30; 72°C-5mins	
Nest 2	rOVA1 TGTAGTATTCAAACGCAGT	800bp	
	rOVA2 TATGTACTTGTTAAGCCTTT	94°C-3mins;{95°C-30s;49°C-30s; 72°C-90s}X30; 72°C-5mins	

Table 2. Constituents of reaction mixture for Nest 1 and Nest 2 reactions of *P. falciparum*, *P. malariae*, *P. vivax* and *P. ovale*

Nest 1		Nest 2	
Reagent	X1 (µl)	Reagent	X1 (µl)
Double-distilled H ₂ O	8.0	Double-distilled H ₂ O	8.25
Pre-mix with BSA (X1)	2.5	Pre-mix with BSA (X1)	2.5
rPLU5 (0.4µM)	0.5	rFAL1/rMAL1/rVIV1/rOVA1 (0.3µM)	0.375
rPLU6 (0.4µM)	0.5	rFAL2/rMAL2/rVIV2/rOVA2 (0.3µM)	0.375
DNA template	1.0	Nest 1 product	1.0
Total	12.5		12.5




Fig. 2. Map of the Adamawa State showing Malaria overall percentage infection by Zones
Source: Department of Geography, University of Port Harcourt

The electrophoresis was allowed to run for 45 minutes and the gel was later viewed under UV light using gel documentation machine.

3. RESULTS

The result in Table 3 shows infection in study locations (hospitals) with their respective Local government areas; the highest infection was recorded in Yola North at Specialist hospital 68.8% with statewide infection of 4.58% followed by Mubi South at General hospital Mubi 57.5% with statewide infection of 3.83% while the least infection was recorded at General hospital Ganye 22.5% with statewide infection of 1.5%. In terms of Parasite density; Yola North had 208896/ μ l followed by Michika 174808/ μ l and Mubi-North 139360/ μ l while the least's parasite density was recorded in Hong 59080/ μ l and Mubi-South 9000/ μ l. Statistical difference for number infected and overall percentage infected $P>0.05$ whereas parasite density $P=.05$.

Table 4 and figure 2 shows infection by zone; in the North it was 45.5% with statewide 15.17%, in the central it was 39.25% with statewide 13.08% and south 32.5% with statewide 10.83%. The difference in infection was statistically significant as $P=.05$. Highest parasite density was recorded in the Central 569828/ μ l followed by South 539572/ μ l and least in the North 523648/ μ l.

3.1 Molecular Analysis

Out of the 108 blood samples examined by molecular analysis 17 were infected with *P. falciparum* as the only species found in Adamawa state by this research. The fragment size detection of 205bp shows the presence of *P. falciparum* (plate 1). Age group 0-5 years old had the highest infection with 26.1% followed by age groups 16-20 years with 22.2% and 6-10 years 18.8% $P=.05$ Table 5. In relation to sex, female was higher with 17.9% infection and male had 13.5% $P=.05$. The total infection was 15.7% Table 6.




Plate 1. Electrophoresis after Plasmodium-specific PCR amplification: L= DNA ladder (100 base pair), P=Positive control; N=Negative control. Lanes 52, 29, 66, 49, 13, 37, and 48 are positive

Table 3. Plasmodium Falciparum Malaria Infection by Locations in Adamawa State

S/No	Locations	No. examined	No. infected (%)	statewide % infected	PD(μ l)	Coordinates
1.	CH Gulak Madagali LGA.	80	22(27.5)	1.83	75656	10°48'17.8"N 13°27'04.5"E
2.	GH Maiha Maiha LGA.	80	39(48.8)	3.25	124824	10°03'54.3"N 13°11'22.3"E
3.	GH Michika Michika LGA.	80	31(38.8)	2.58	174808	10°38'29.9"N 13°23'03.9"E
4.	NMLC Mubi Mubi North LGA.	80	44(55)	3.67	139360	10°16'07.8"N 13°17'02.2"E
5.	GH Mubi Mubi South LGA.	80	46(57.5)	3.83	9000	10°15'43.7"N 13°16'12.2"E
6.	CH Fufore Fufore LGA.	80	29(36.3)	2.42	138728	9°12'56.55204"N 12°39'49.63896"E
7.	GH Garkida Gombi LGA.	80	25(31.3)	2.08	63996	10°24'40.8"N 12°33'56.7"E
8.	CH Hong Hong LGA.	80	28(35)	2.33	59080	10°14'06.0"N 12°56'44.3"E
9.	CH Song Song LGA.	80	20(25)	1.67	99128	9°49'10.0"N 12°37'15.2"E
10.	SH Yola Yola North LGA.	80	55(68.8)	4.58	208896	9°16'40.6"N 12°26'47.6"E
11.	GH Borrong Demsa LGA.	80	35(43.8)	2.92	88428	9°32'29.4"N 12°10'23.3"E
12.	GH Ganye Ganye LGA.	80	18(22.5)	1.5	95456	8°26'10.1"N 12°03'47.1"E
13.	GH Numan Numan LGA.	80	28(35)	2.33	138068	9°28'06.6"N 12°01'56.1"E
14.	CH Mayo-Belwa Mayo-Belwa LGA.	80	25(31.3)	2.08	105716	9°03'34.7"N 12°03'52.2"E
15.	CH Toungo Toungo LGA.	80	24(30)	2.0	111904	8°07'23.9"N 12°02'48.4"E
Total		1200	469(39.08)		1633048	

GH= General hospital, CH= Cottage hospital, SH= Specialist hospital PD= Parasite density, S/No= Serial number, %=percentage. No positive and % infected: $P>0.05$, PD: $P=.05$.

Table 4. Malaria infection by geographical zone in Adamawa state

S/No	Zone	No. examined	No. infected	% Infected	Statewide % infected	PD(μ l)
1.	North	400	182	45.5	15.17	523648
2.	Central	400	157	39.3	13.08	569828
3.	South	400	130	32.5	10.83	539572
Total		1200	469	39.08	39.08	1633048

PD= Parasite density, μ l= microliter, $\chi^2=14.204$, $P=0.001$

The Adamawa State overall data results in relation to age groups shows age 0-5 years old had the highest infection with 46.2% followed by 6-10 years with 42.5% and the least was recorded in age group 16-20 years 35.4% $P=.05$. In terms of parasite density age group 41 and above year's old recorded 321600/ μ l while the least was recorded within age group 6-10 years 100512/ μ l Table 7.

In relation to sex, females had 40.3% infection with parasite density of 967252/ μ l and males

recorded 37.3% infection with parasite density 665796/ μ l $P=.05$ Table 8.

According to socioeconomic class, the middle class had the highest infection with 43% followed by the higher class 38.6% whereas the lower class recorded 38.1% infection $P=.05$. The highest parasite density was recorded in lower class 926603/ μ l and least in middle class with 338605/ μ l Table 9.

Table 5. Plasmodium falciparum malaria detected by molecular identification in relation to age in Adamawa State

S/No.	Age group(years)	No. examined	No. infected	% infected	P-value
1.	0-5	23	6	26.1	0.000
2.	6-10	16	3	18.8	
3.	11-15	6	0	0	
4.	16-20	9	2	22.2	
5.	21-25	20	2	10	
6.	26-30	11	2	18.2	
7.	31-35	6	0	0	
8.	36-40	6	1	16.7	
9.	41&above	11	1	9.1	
Total	108	17	15.7		

S/No = serial number, % = percentage

Table 6. Plasmodium falciparum malaria detected by molecular identification in relation to sex in Adamawa State

S/No.	Sex	No. examined	No. infected	% infected	P-value
1.	Female	56	10	17.9	0.021
2.	Male	52	7	13.5	
	Total	108	17	15.7	

S/No = serial number, % = percentage

Investigating malaria parasitaemia in relation to possession and frequency of sleeping under mosquito net, it was found that subjects who do not have mosquito net recorded highest infection 52.3% followed by those who do not sleep under mosquito net (because is stuffy) 50% whereas those reported to sleep under

mosquito net (non-ITN) regularly recorded 46.7% infection $P=0.05$. Those who reported to sleep under treated mosquito net regularly recorded 658708/ μ l parasite density and the 4 infected non-regular users of untreated mosquito net recorded least parasite density 12600/ μ l Table 10.

Table 7. Age related malaria parasitaemia in Adamawa State

S/No	Age group(yrs)	No examined	No infected	% infected	Parasite density(μ l)	P-value
1.	0-5	119	55	46.2	189184	0.000
2.	6-10	80	34	42.5	100512	
3.	11-15	83	33	39.8	112472	
4.	16-20	144	51	35.4	162564	
5.	21-25	138	56	40.6	184612	
6.	26-30	120	49	40.8	167236	
7.	31-35	149	53	35.6	188176	
8.	36-40	165	61	40	206692	
9.	41 & above	202	77	38.1	321600	
	Total	1200	469	39.1	1633048	

S/No= Serial number, yrs= years, %= percentage, μ l= microliter**Table 8. Malaria parasitaemia in relation to sex in Adamawa State**

S/No	Sex	No examined	No infected	% infected	Parasite density(μ l)	P-value
1.	Female	660	266	40.3	967252	0.000
2.	Male	540	203	37.6	665796	
	Total	1200	469	39.1	1633048	

S/No= serial number, %= percentage, μ l= microliter

Table 9. Malaria parasitaemia in relation to socioeconomic status in Adamawa State

S/No	Socioeconomic class	No examined	No infected	% infected	Parasite density(μ l)	P-value
1.	Higher class	293	113	38.6	367840	0.000
2.	Middle class	221	95	43	338605	
3.	Lower class	686	261	38.1	926603	
	Total	1200	469	39.1	1633048	

%=percentage, μ l=microliter

Table 10. Malaria parasitaemia in relation to possession and frequency of sleeping under mosquito net in Adamawa State

S/No	Possession and frequency of sleeping under mosquito net.	No Examined	No infected	% infected	P.D(μ l)	P-value
1.	Do not have mosquito net	132	69	52.3	272964	0.000
2.	Regular sleeping under mosquito net (ITN)	578	201	34.8	658708	
3.	Regular sleeping under mosquito net (non-ITN)	15	7	46.7	34332	
4.	Non-regular sleeping under Mosquito net (ITN)	451	183	40.6	621668	
5.	Non-regular sleeping under Mosquito net (non-ITN)	14	4	28.6	12600	
6.	Do not sleep under mosquito net (is stuffy)	10	5	50	32776	
	Total	1200	469	39.1	1633048	

S/No= serial number, %= percentage, P.D= parasite density, %= percentage, μ l= micrometer

4. DISCUSSION

This study investigated the prevalence of malaria in relation to socioeconomic status, treatment seeking behavior, mode of prevention practiced; in particular the use of mosquito nets in Adamawa state. In this study the only species of Plasmodium found by both microscopy and molecular analysis was *P. falciparum* which agrees with report that falciparum malaria is responsible for 99.7% of estimated malaria cases in Africa [22]. It also concurs with findings in Rivers state, Nigeria [23] and Kano, Nigeria [24]. Out of the total 1200 samples examined by microscopy 469 (39.1%) were infected, the total parasite density was 1633048/ μ l, this is of utmost concern as reports also indicate the rise in malaria cases in Adamawa state [7].

By location Yola North recorded highest infection 68.75% with overall state infection of 4.58% and parasite density 208896/ μ l. This finding is slightly higher than reported 50.6% infection among children in Yola, but, agrees with reports that Yola North had the highest malaria cases [25]. The high infection rate in Yola North could be attributed to, warm temperature, densely populated settlements and proximity to River

Benue along which both rainy and dry season farming activities takes place; hence favourable conditions for Anopheles mosquitoes to breed and transmit infection at ease. Other factors could be lack of compliance to regular sleeping under mosquito net because of hot weather condition. Mubi South and Mubi North recorded 57.5% (with statewide 3.83%) infection and 55% (with statewide 3.66%) infection respectively. The presence of River and traditional dug well in houses and communities (as observed during field work) could serve as contact sites with mosquitoes amidst other factors like clustered city setting leading to high rate of infection in Mubi town. In addition, the indiscriminate dumping of refuse could serve as breeding sites for mosquito in Mubi town [26]. Difference in number of infection by location in Adamawa state was not statistically significant as $P > 0.05$ whereas parasite density was significant as $P = 0.05$. Infection by zones showed Adamawa North recorded 45.5% with statewide infection of 15.17% and total parasite density of 523648/ μ l this could not be unconnected with the presence of Anopheles mosquitoes [17] in addition to predominantly farming occupation. Most communities still use dug well even in houses that are without window/door nets hence

exposing locals to mosquito bite before and after bed time (i.e if any mosquito net). Adamawa central recorded 39.25% infection with state wide infection of 13.08% and parasite density 569828/ μ l. This is not surprising because the breeding sites of mosquito along River Benue are very possible; rice and vegetable farmers, fishermen and pastoralists could easily be exposed to infection. Less infection was recorded in most parts of Adamawa South with exceptions to LGAs along the Benue River (Demsu and Numan); Adamawa south recorded 32.5% infection with statewide infection 10.83% and parasite density 539572/ μ l. A relative low malaria cases in Adamawa South was reported, especially around Toungo LGA and Ganye LGA possibly due to proximity to Gashaka Gumti National park which has low temperature and dense forest making it unfavourable for mosquito breeding [27]. The difference in percentage infection by zone was significant as $P=.05$.

The overall data obtained from Adamawa state in relation to age showed, the highest infection was recorded within age group 0-5 years old 46.2% with parasite density 189184/ μ l, same with molecular analysis 0-5 years recorded highest infection with 26.1% this concur with reports that most malaria deaths occur in children less than 5 years as they are among the vulnerable groups to infection due to low immunity [7,28]. The least infection was recorded among age group 16-20 years 34.5% with parasite density 162564/ μ l. The less infection could be due to acquired immunity; this finding is higher than the 26% infection reported in age group 11-21 years [29]. Infection by age groups was statistically significant as $P=.05$.

In relation to sex, females recorded higher infection 40.3% and males had 37.6% infection $P=.05$. Parasite density by sex was females 967252/ μ l and males 665796/ μ l. Molecular analysis result also showed higher infection in females 17.9%, than males 13.5% $P=.05$. The higher infection in females may be gender related as women are known to engage in outdoor house chores at dawn and dusk a favourable moment for vector mosquito bite. Higher malaria infection in females than males was also reported in Yola [25] and in Emohua, Rivers state, Nigeria [23].

According to socioeconomic status, the middle class recorded highest infection 43% followed by higher class 38.6% $P=.05$. The lower class recorded highest parasite density 926603/ μ l

followed by higher class 367840/ μ l. It was reported that high burden of malaria is directly related to poverty [22,30] which concurs with this finding in terms of parasite density.

Further study was carried out on infection in relation to possession and sleeping under mosquito net. It was found that subjects who do not have mosquito net recorded 52.3% infection followed by subjects who do not sleep under mosquito net (because is stuffy) with 50% infection whereas those who sleep under mosquito net (ITN) regularly had 34.8% infection. Access and ownership of mosquito net does not translate to its use, as revealed in this study, some findings reports the misuse of mosquito nets for other purposes such as fishing [31], hence ownership is not enough [32]. There was a high parasite density 658708/ μ l among subjects who sleep under mosquito net which could be as a result of contact with mosquitos outdoors. There was statistical difference in infection related to possession and ownership of mosquito net as $P=.05$.

5. CONCLUSION

Considering the high infection of malaria in Adamawa state 39.1%, there is need for sustained interventions with insecticidal treated mosquito nets and chemoprevention, with rate of infection by locations in mind. Sustained awareness on importance of sleeping under mosquito net should be encouraged as possessing it does not necessarily translate into proper use. Other methods of prevention such as environmental sanitation and wearing protective clothing are encouraged while awaiting the vaccine roll-out for children as additional tool. Free malaria communities could lead to improved socioeconomic status as the infection is associated with economic difficulty.

CONSENT AND ETHICAL APPROVAL

Clearance was obtained from University of Port Harcourt Research ethics committee in 2019 and Adamawa State Ministry of Health Research Ethical Committee in 2021 with approval number ADHREC 5/03/2021/043. All subjects and guardian of children under 18 years gave written informed consent to participate in providing blood sample. A verbal explanation of the objectives, benefits (such as interest groups' possible intervention to eliminate malaria which could improve standard of living in the study area) and potential risks associated with participation in the

study were given to all subjects. Subjects' personal information was kept confidential.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

ACKNOWLEDGEMENT

We thank the Research Ethics committee of the University of Port Harcourt and Adamawa State Ministry of Health Research Ethical Committee for approving this work. We are grateful to all the laboratory staff of the selected secondary health facilities for their kind support during this study. We also appreciate all subjects who gave us their consent for sample collection.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. <https://stanfordhealthcare.org/medical-conditions/primary-care/malaria/types.html>
2. Enock K, Pukuma SM, Augustine LM, Gundiri L.B., Zamdayu NM and Daniel JL. Prevalence and associated risk factors of malaria infection among pregnant women attending antenatal clinic in Yola North, Adamawa state. *The zoologist* 2002;18:19-25. Available:<https://dx.doi.org/10.4314/tzool.v18i1.4>
3. Warral E, Suprotik B, Hara H. The relationship between socioeconomic status and malaria: A review of the literature; 2003. Available:https://assets.publishing.service.gov.uk/media/57a08c12e5274a27b2000f71/WP01_03.pdf.
4. Somi MF, Kachur, SP, Butler J, Njau JD, Vahid, F, and Abdulla S. Is there evidence for dual causation between malaria and socioeconomic status? Findings from rural Tanzania. *The American journal of tropical medicine and hygiene.* 2007;77 6:1020-7.
5. World Health Organisation. Malaria keyfacts; 2023. Available:<https://www.who.int/news-room/fact-sheets/detail/malaria>
6. World Health Organisation. Malaria fact sheet, 2018.
7. World Malaria report 2022. World Health Organisation 2022. Licence:CC BY-NY-SA3.0 IGO.
8. Nigeria Malaria Fact Sheet. United State Embassy in Nigeria; 2011.
9. Sokunbi TO, Omojuyigbe JO, Bakenne HA. and Adebisi YA. (2022). Nigeria End Malaria Council: What to expect. *Annals of medicine and surgery* (2012);82:104690. Available:<https://doi.org/10.1016/j.amsu.2022.104690>
10. Available:<https://reliefweb.int/report/nigeria/mothers-adopt-seasonal-malaria-chemoprevention-drugs-keep-malaria-bay-adamawa-state>
11. Available:<https://www.who.int/news/item/05-07-2023-18-million-doses-of-first-ever-malaria-vaccine-allocated-to-12-african-countries-for-2023-2025--gavi--who-and-unicef>
12. Available:<https://www.reuters.com/world/africa/nigeria-regulator-grants-approval-oxfords-malaria-vaccine-2023-04-17/>
13. Available:https://www.citypopulation.de/en/nigeria/admin/NGA002_adamawa/
14. Available:<https://adamawastate.gov.ng/history/>
15. Available:https://en.wikipedia.org/wiki/Adamawa_State
16. Available:<https://itsmyschoollibrary.com/2020/03/07/adamawa-state-history-topography-occupation-tourism-festivals-local-government/>
17. Wahedi JA., Ande AT., Oduola AO, Obembe A, Tola M, Oyeniyi TA and Awolola TS. Dynamics of malaria vector indices in two vegetation zones within North Eastern Adamawa state, Nigeria. *Tropical Biomedicine.* 2020;37(3): 637-649.
18. Glenn DI. Determining sample size. *Programme evaluation development, Florida PEOD-6* 2003:4.
19. www.mobile-CDR.com
20. World Health Organisation. Malaria parasite counting. Malaria microscopy standard operating procedure-MM-SOP-09; 2016.
21. Snounou G, Viriyakboso S, Zhu XP, Jarra W, Pinheiro L, doRosario VE, et al. High sensitivity of detection of human malaria parasites by the use of nested polymerase chain reaction. *Mol Biochem Parasitol* 1993; 61:315-320.
22. The World 'malaria report at glance' 2019.

- Available:<https://www.who.int/news-room/feature-stories/detail/world-malaria-report-2019>
23. Richard A, Nduka FO, Awi-waadu GDB., and Wogu MN. Malaria parasitaemia among selected residents of Emohua Local Government Area and Port Harcourt metropolis, Rivers state, Nigeria. *Nigerian Journal of Parasitology*. 2019;4(1): 32-36.
 24. Salwa D, Hesham M, Al-Mekhlafi, II, Jamaiah I, Wahid MA, Awatif MA *et al*. Is Nigeria winning the battle against malaria? Prevalence risk factors and KAP assessment among Hausa communities in Kano state, *Malaria Journal*. 2016;15. DOI:10.1186/s12936-016-1394-3.
 25. Kunihya IZ, Samaila AB, Nassai I, Sarki A, and Haruna MY. Prevalence of malaria infection among children attending specialist hospital Yola, Adamawa state, Nigeria. *Journal of Medical and Biological Science Research*. 2016;1(8): 132-142.
 26. Jummai VZ, Garandi ID, and Awal Y. An assessment of trends and spartial distribution of cases of malaria fever in Mubi South Local Government Areas, Adamawa State, Nigeria. *Journal of Environmental & Material Sciences*. 2021;1:2:72-82.
 27. Kukur JH, Elishama LJ, Joshua DZ, John AG and Adamu BM. Regional trends of malaria cases in Adamawa state. *Global Scientific Journals*. 2019;8:732-737.
 28. Abossie A, Yohanes T, Nedu A, Tafesse W and Damitie M. Prevalence of Malaria and Associated Risk factors Among Febrile children under five years: A cross-sectional study in Arba Minch Zuria District, South Ethiopia. *Infection and drug resistance*. 2022;13:363-372. Available:<https://doi.org/10.2147/IDR.S223873>
 29. Naphtali RS, Babylon P, Musa ZJ, Atinga A. Prevalence of Malaria Infection in Yola-South L.G.A. of Adamawa state. *IORS Journal of Pharmacy and Biological sciences* 2017; 12:5:58-61.
 30. Health think team. *Malaria in Nigeria: Where are we now?*; 2017.
 31. Bertozzi-Villa A, Bever CA, Koenker H, Weiss DJ, Vargas-Ruiz C, Nandi AK, *et al*. Maps and metrics of insecticide-treated net access, use and nets-per-capita in Africa from 2000-2020. *Nature communications* 2021;12: 3589.
 32. Orish VN, Maalman RSE, Donkor OY, Cemantis BYM, Osei E, Amu H, *et al*. Assessing health-seeking behavior and malaria prevention practices among communities in four districts of the Volta region of Ghana. *Malaria Journal* 2021; 20:450.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/117562>