



Phytochemical Screening and Evaluation of the Cytotoxicity of Fruits of *Solanum torvum* Swartz (Solanaceae) on HFF Cells (Human Foreskin Fibroblasts)

Kanga Yao^{1*}, Tra Bi Irié Otis², Ambe Alain Serge³, Camara Djeneb¹ and Zirihi Guédé Noël¹

¹Botanical Laboratory, UFR Biosciences, Felix Houphouët Boigny University, P.O.Box 582, Abidjan 22, Côte d'Ivoire.

²Biochemical Microbiological Laboratory of SVT, Higher Teacher Training School of Côte d'Ivoire, P.O.Box 10, Abidjan 08, Côte d'Ivoire.

³Botany Laboratory of SVT, Higher Teacher Training School of Côte d'Ivoire, P.O.Box 10, Abidjan 08, Côte d'Ivoire.

Authors' contributions

This work was carried out in collaboration between all authors. Author KY designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors TBIO and AAS managed the analyses of the study. Authors CD and ZGN managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRIB/2018/43993

Editor(s):

(1) Dr. K.S Vinayaka , Assitant Professor, Department of Botany, Kumadvathi First Grade College, Shimoga, Karnataka, India.

Reviewers:

(1) Esraa Ashraf Ahmed ElHawary, Ain Shams University, Egypt.

(2) Uchendu, Mbah Okuwa, Michael Okpara University of Agriculture, Nigeria.

(3) Jean-De-Dieu Tamokou, University of Dschang, Cameroon.

(4) R. Dhivya, Nirmla College for Women, India.

Complete Peer review History: <http://www.sciencedomain.org/review-history/26685>

Original Research Article

Received 18th August 2018
Accepted 10th October 2018
Published 19th October 2018

ABSTRACT

Objective: The aim of this work is to evaluate the cytotoxic activity of the 70% ethanolic extract of the fruits of *Solanum torvum* Swartz (Solanaceae) on HFF (Human Foreskin Fibroblast) cells in *in vitro* culture and to determine its phytochemical composition.

Study Plan: An ethnobotanical survey was conducted in August 2015 in the Haut-Sassandra

Region (Ivory Coast), on medicinal plants with multiple uses, and at the end of this survey the fruits of *Solanum torvum* have been selected then harvested in the Sub-prefecture of Bédiala (Ivory Coast). Then after drying, the ethanolic extract of the fruits was prepared and sent to France in February 2016 at the Laboratory Adaptation and Pathogenesis of Microorganisms (LAPM) of Grenoble for cytotoxic tests. Phytochemical screening was carried out at the Faculty of Biological Sciences of Félix Houphouët Boigny University (Côte d'Ivoire).

Methods: From an ethnobotanical survey, the fruits of *Solanum torvum* were harvested. The 70% ethanolic extract prepared from the fruits of this plant was tested *in vitro* on divisional HFF cells after phytochemical screening.

Results: The result revealed that this extract has cytotoxic activity on the tested HFF cells. At 800 µg/mL, the survival rate of HFF cells increased from 100% to 4% of living cells. Phytochemical screening revealed the presence of compounds such as alkaloids, tannins, polyphenols, saponins and flavonoids.

Conclusion: This extract is cytotoxic on HFF cells. It is, therefore, necessary to continue studies on toxicity and to be cautious in the use of *solanum torvum* fruits in traditional medicine.

Keywords: HFF cell; cytotoxic; extracts; ethnobotanical survey; *Solanum torvum*; phytochemicals.

1. INTRODUCTION

Solanum torvum belongs to the family Solanaceae. *Solanum torvum* is native to Central and South America, from Mexico to Brazil and Peru. It has spread widely in the Caribbean [1]. *Solanum torvum* belongs to the family Solanaceae. In West and Central Africa, it is grown locally in gardens for cooking [2]. Fruits are widely used in the treatment of shingles in Cameroon [3]. They are also used as a vegetable and considered an essential ingredient in the diet of the South Indian population [1]. A fruit decoction is used in Ghana for the treatment of cough, liver disease and spleen [4]. *Solanum torvum* are rich in antioxidant, Its extracts are used in the preparation of tonic and hemopoietic agents and also for the treatment of pain throughout the body [5,6]. Previous work has revealed that the fruits of *Solanum torvum* are used in the treatment of various conditions: microbial infections [7], hypertension [8], kidney disease [9] diabetes [10]. Pérez-Amador et al. [11] reported the presence of glycoalkaloids in the fruit of *Solanum torvum* and another study showed that very low doses of glycoalkaloids in some Solanaceae are toxic [12,13]. In view of the interest of *Solanum torvum* fruits in the traditional medicine, we have undertaken to evaluate scientifically the cytotoxicity and phytochemical screening of the fruits of *Solanum torvum*. The aim of this work is to perform a phytochemical screening and a cytotoxicity study of *Solanum torvum* fruits on Human Foreskin Fibroblasts (HFF), cells that intervene in the anti-infectious defence of an organism.

2. MATERIALS AND METHODS

2.1 Plant Material

The fruits of *Solanum torvum* (Fig. 1) were harvested in Biadiala in the Department of Daloa (Ivory Coast).

2.2 Cellular Material

The cellular support consists of human HFF (Human Foreskin Fibroblasts) cells. These are human cells that testify to the toxic activity of an extract. When these cells are in culture for only 24 hours, they are in a state of mitosis (or dividing cells).

2.3 Preparation of Plant Extracts

After harvest, the fruits were freed of impurities, dried in the shade for a week and then pulverised with an electric grinder. The fine powders obtained were stored in glass jars to prevent mould.

2.4 Preparation of Total Aqueous Extract (TAE)

The preparation of this extracts was performed using the method described by Zirih et al. [14] which involves macerating 100 g of plant powder of species in 1L of sterile distilled water using a Blender type 7 SEVEN STAR. The homogenate was filtered over hydrophilic cotton and then on Whatman filter paper (n°3). The aqueous filtrate thus obtained is evaporated in an oven of Med Center Venticell type at 50°C to obtain powders that constitute the total aqueous extract (TAE).



Fig. 1. Leafy and fruiting twig of *Solanum torvum* (Solanaceae)

2.5 Preparation of 70% Ethanolic Extract (70% FE)

The extract was obtained by dissolving 5 g of TAE in 100 mL of 70% ethanolic (67.2 mL of pure 96% ethanolic for 28.8 mL of distilled water) solution and then homogenised. After decantation and filtration of the alcoholic fraction on hydrophilic cotton and on Whatman filter paper (n°_3), the filtrate collected is evaporated in an oven at 50°C. The powder obtained constitutes 70% ethanolic extract (70% EE) [15].

2.6 Yield Calculation

The yield is the quantity of extract obtained from the plant powder. It is expressed as a percentage. In practice, it has been determined by the ratio of the weight of the solids content after evaporation on the weight of the dry plant material powder used for the extraction, multiplied by 100. This result is indicated by the following formula:

$$Yd = (m \times 100)/M$$

Yd : Extraction yield in percentage
 m : mass in grams of the dry extract
 M : mass in grams of the drug powder.

2.7 Phytochemical Screening

The identification of different chemical compounds in 70 % ethanolic was done by tubes characterisation reactions. This method consists of detecting the different families of chemical compounds that may exist in plant extracts on

the basis of characteristic colourations or precipitation reactions [16].

2.7.1 Alkaloids characterisation

The characterisation of alkaloids was made using Bouchard (iodo-iodide) and Dragendorff (tetraiodo potassium bismuthate) reagent. 6 mL of 70% ethanolic extract solution was evaporated to dryness. The residue was taken up in 6 mL of alcohol at 60°C. The filtrate thus obtained was divided into two test tubes. In the first tube, two drops Dragendorff reagent were added. The presence of alkaloids was characterised by observing orange-coloured precipitates. In the second tube, two drops of Bouchard reagents was added. The appearance of a reddish-brown colour indicates the presence of alkaloids. A control test was made with quinine.

2.7.2 Characterisation of polyphenols

The polyphenols colorimetry forms coloured precipitate with a solution of ferric chloride ($FeCl_3$). Thus, one drop of alcoholic solution of 2% ferric chloride and 2 mL of solution of 70 % ethanolic extract was added. The formation of blue-black or green colouring more or less dark confirms to the presence of polyphenols. A control test was performed with a solution of phenol.

2.7.3 Characterisation of flavonoids

Flavonoids have been characterised by the reaction to cyanidin. Thus, 2 mL of 70% ethanolic extract were evaporated to the dry sand bath.

The residue thus obtained was mixed with 5 mL dilute hydrochloric acid 2 times. The mixture was collected in a test tube, in which pink-orange or violet colouration will appear. The addition of 3 drops of isoamyl alcohol intensifies this colouring and confirms the presence of flavonoids. An alcoholic solution of quercetin was used as a control.

2.7.4 Tannins characterisation

The Stiasny reagent (Formalin 30%, concentrated HCl 1/0.5) helped to distinguish the catechin tannins (by precipitation) of gallic tannins (by saturation). Tannins cathéchiqes: to 10 mg of 70% ethanolic extract, were added 10 mL of Stiasny reagent. The mixture was heated in a water bath at 80°C for 30 minutes. After cooling in a stream of water, observation of precipitate in the form of clear-brown flakes characterises catechin tannins. An alcoholic solution of catechin was used as a control. Gallic tannins: For this test, the filtrate obtained from the reaction of catechol tannins characterisation was saturated with sodium acetate. To this mixture was added a few drops of a dilute aqueous solution of FeCl₃ at 1% (approximately 1 mL). The appearance of an intense blue-black colouration indicates the presence of gallic tannins not precipitated by Stiasny reagent. An alcoholic solution of gallic acid was used as a control.

2.7.5 Terpenes characterisation

Sterols and terpenes characterisation was made by the Liebermann-Burchard reaction. To 0.2 g of 70 % ethanolic extract, were added 5 mL of ethyl ether, then the mixture was macerated for 30 minutes. The solution obtained after the maceration was filtered and then evaporated to dryness. The residue was then dissolved in 0.5 mL of acetic anhydride. Using a pipette, 2 mL of concentrated sulfuric acid were laid down at the bottom of the test tube without stirring. The appearance of brownish red or purple ring reflects the two liquid contact zone. The upper liquid turns green or purple to green or purple indicating the presence of sterols and terpenes. A control test was performed with progesterone.

2.7.6 Coumarins characterisation

For the detection of coumarins, 2 mg of 70 % ethanolic extract was added to 2 mL of warm water and then homogenised. The homogenate thus obtained was divided into two test tubes.

There after, 0.5 mL of diluted ammonia at 25% was added to the contents of one of the tubes. After observation under UV 365 nm, the presence of fluorescence in the tube where ammoniac was added indicates the presence of coumarins.

2.7.7 Saponins characterisation

For the detection of saponins, 10 mL of 70% ethanolic extract was introduced in the test tubes. Each tube was strongly stirred in a vertical position for 15 seconds, and then left to settle 15 minutes. The height of persistent foam is higher than 1 cm, testifying the presence of saponins.

2.8 Cytotoxicity Test

To measure the toxicity of the ethanolic extract, the Human Foreskin Fibroblasts (HFF) cells were inoculated in 96-well plates (CellStar) at the rate of 3000 to 5000 cells per well in 100 µl of D10 medium. These cells are kept in culture for 24 hours (dividing cells). Subsequently they were exposed for 24 hours at different concentrations (125-800 µg /mL) in plant extract solubilised in PBS buffer. This was done in triplicate, also for control control without plant extract. Viability was determined using 3- (4, 5-dimethylthiazol-2-yl) - 2,5-diphenyl tetrazolium bromide (MTT). The tetrazolium ring it contains is reduced in formazan by the mitochondrial succinate dehydrogenase of metabolically active cells, which precipitates and gives a purple colour. The amount of precipitate formed is proportional to the number of living cells. In each well, MTT is added at a concentration of 500 µg / mL and incubated for 3 hours at 37 ° C. The formazan crystals are solubilised in 10 mM dimethylsulfoxide (DMSO). The measurement of the optical density at 544 nm was made using a Safir spectrophotometer (Tecan); this measurement of absorbance will make it possible to determine the relative quantity of living and metabolically active cells [17]. The results were expressed as a percentage of viability compared to control without plant extract. Viability rate = (Abs544 nm extract / Abs544 nm control) × 100.

3. RESULTS

3.1 Yield of Different Extracts of Fruits of *Solanum torvum*

We obtained from 200 g of powder, 20 g of total aqueous extract a yield of 10% and from 5 g of

total aqueous extract, we got 2 g of 70% ethanolic extract or a yield of 40%.

3.2 Phytochemical Sorting

The phytochemical sorting performed with the extracts of fruits of *Solanum torvum* allowed to detect the presence of various chemical groups (Table 1). They are the polyphenols, tannins, flavonoids, saponins, and alkaloids in 70% ethanol extract.

3.3 Cytotoxicity Test

Fig. 2 gives the percentage of viability of the HFF cells cultured in the presence of concentrations of 100 to 800 µg / mL for the 70% ethanolic extract of the fruits of *Solanum torvum* compared to the control without plant extract. The number of cells decreases considerably as the

concentration of the 70% ethanol extract of the fruits of *Solanum torvum* increases. At 800 µg / mL the number of dividing cells is 4%. The averages with the same superscript letters are not different at 5% according to the Turkey test.

4. DISCUSSION

Medicinal plants play a central role in traditional medicine. Ethnobotanical surveys conducted among traditional health practitioners have made it possible to harvest the fruits of *Solanum torvum*, which is used in the treatment of anaemia, bacterial infections and several other diseases [18]. The recipes obtained from the fruits of this plant are monospecific, which is an advantage for the patients, because the associations of wrongly mixed plants, are sometimes dangerous for the health [19].

Table 1. Chemical compounds in the fruits of *Solanum torvum*

Species	Extract	Chemical compounds							
		Sap	Flav	Terp/ster	Tanins		Coum	Alc	Poly
					Gall	Cathé			
<i>Solanum torvum</i>	EE 70%	+	+++	-	++	++	-	+	+

- : negative reaction ; + : positive reaction; EE 70% : 70% ethanolic extract; Sap: Saponins; Flav: Flavonoids; Terp / Ster: Terpenes / sterols; Gall: Gallic; Cathé: Catechic; Coum: Coumarines; Alc: Alkaloids; Poly: Polyphenol




Fig. 2. Cytotoxicity test of 70% ethanolic extract of fruits of *Solanum torvum* on HFF dividing cells. Data expressed as mean ± ecart-type (n=3)

Phytochemical screening revealed the presence of chemical compounds such as alkaloids, tannins, polyphenols, saponins and flavonoids. The presence of these chemical compounds could justify the multiple activities of the fruits of this plant [20]. The cytotoxic assay performed on HFF cells showed a gradual decrease in purple staining in each well. Since the dye penetrates only in living cells, the colouring is weaker as the plant extract is cytotoxic by inhibition of HFF cells [21]. The sharp decrease in the relative amount of the dividing HFF cells could be explained by the fact that the HFF cells would be killed by the 70% ethanol extract of *Solanum torvum*. Indeed, extracts resulting in a cell death greater than 30% could be considered as cytotoxic [22]. This extract could therefore contain a chemical compound that inactivates succinate dehydrogenase, an enzyme important for mitochondrial respiration, the blockage of which would lead to cell death. This result demonstrates the cytotoxic effect of 70% ethanolic extract of *Solanum torvum*, a Solanaceae from the Ivorian pharmacopoeia on the cell line tested. Which means that the external use of the fruits of this plant would probably be dangerous for human health. This toxicity of fruit could also be explained by the presence of certain groups of chemical compounds such as glycoalkaloids which are toxic in some Solanaceae [13]. Our results on *in vitro* toxicity corroborate those [23] who worked on the same family of plants. Indeed according to the work of Busser and Baies [23] the fruits of *Solanum nigrum* L. (Solanaceae) another Solanaceae rich in glycoalkaloids and saponins are toxic in internal and external uses on an organism.

5. CONCLUSION

Solanum torvum is an important medicinal plant of the family of Solanaceae. From the evaluation of the biological activity, it appears that 70% ethanolic extract of the fruits of *Solanum torvum* is cytotoxic on the HFF cells. Hence, further studies on the cytotoxic activities of this plant extract is warranted.

ACKNOWLEDGEMENT

The authors thank the Laboratory Adaptation and Pathogenesis of Microorganisms (LAPM) of Grenoble in France, where cytotoxicity studies were carried out and the traditional health practitioners of the Haut-Sassandra Region.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Adjanohoun J, Aboubakar N, Dramane K, Ebot E, Ekpere A, Enoworock G, et al. Traditional medicine and pharmacopoeia-contribution to ethnobotanical and floristic studies in Cameroon. In: CNPMS. Porto-Novo, Benin. 1996;50–52.
2. Etienne J. Solanacées médicinales et philatélie. Bull. Soc. Pharm. Bordeaux. 2005;144:311-332.
3. Emmanuel N, Catherine K, Philomène CNB, Siegfried DD, Emmanuel MM. Inventaire et caractérisation des plantes médicinales utilisées en thérapeutique dans le département de la Sanaga Maritime: Ndom, Ngambe et Pouma. J. Appl. Biosci. 2016;106:10333–10352.
4. Kala CP. Ethnomedicinal botany of the Apatani in the Eastern Himalayan Region of Indian. J Ethno and Ethnomed. 2005;1: 1-8.
5. Sivapriya M, Srinivas L. Isolation and purification of a novel antioxidant protein From the water extract of Sundakai (*Solanum torvum*) seeds. Food Chemistry. 2007;104:510-517.
6. Waghulde H, Kamble S, Patankar P, Jaiswal BS, Pattanayak S, Bhagat C, et al. Antioxidant activity, phenol and flavonoid contents of seeds of *Punica granatum* (Punicaceae) and *Solanum torvum* (Solanaceae). Pharmacologyonl. 2011;1: 193-202.
7. Lalitha V, Raveesha K, Kiran B. Antimicrobial activity of *Solanum torvum* swart. Against Important Seed Borne Pathogens of Paddy, IJEE. 2010;1(2):160-164.
8. Mohan M, Jaiswal BS, Kasture S. Effect of *Solanum torvum* on blood pressure and metabolic alterations in fructose hypertensive rats. J Ethnopharmacol. 2009;126:86–89.
9. Mohan M, Kamble S, Kasture S. Protective effect of *Solanum torvum* on doxorubicin-induced nephrotoxicity in rats. Food and Chem Toxicol. 2010;48:436-440.
10. Gandhi GR, Ignacimuthu S, Paulraj MG, Sasikumar P. Antihyperglycemic activity and antidiabetic effect of methyl caffeate isolated from *Solanum torvum* Swartz.

- Fruit in streptozotocin induced diabetic rats. Eur J Pharmacol. 2011;30:623-31.
11. Pérez-Amador MC, Muñoz OV, García JM, Castañeda AR, González E. Alkaloids in *Solanum torvum* Sw (Solanaceae). Inter experim Botany. 2007;76:39-45.
 12. Tajner-Czopek A. Food Chem. 2008;106 (2):706-11.
 13. Ayad K. Effect of Solanine on arthritis symptoms in postmenopausal female albino rats. Arab Journ of Nucl Sc and Appl. 2013;46(3):279-285.
 14. Zirihi GN, Kra A, Dadié ET. Etude botanique et évaluation des activités antifongiques de *Mitracarpus villosus* (MV) (Rubiaceae) et *Spermacoce verticillata* (SV) (Rubiaceae) sur la croissance *in vitro* de *A. fumigatus*. Rev de Méd et de Pharma Afr. 2007;20:9-17.
 15. Gnahoué G, Bené K, Coulibaly K. Etude botanique, screening phytochimique et activité anticandidosique *in vitro* de *Pycnanthus angolensis* (Welw.) Warb. (Myristicaceae). Euro Sc Jour. 2015;11 (36):7431-7881.
 16. Harborne JB. A guide to modern techniques of plant analysis. Springer, 3rd Edn, India (New Delhi). 1998;5-32.
 17. Mosman T. Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assay. Journal of immunological Methods. 1983;65: 55-63.
 18. Asiedu-Darko E. A survey of indigenous knowledge about food and medicinal properties of *Solanum torvum* in East Akim District of Eastern Region of Ghana. Ghana Jour of Agri Sci. 2010;43(5): 61–64.
 19. N'Guessan K, Kouadio K, Kouamé NF, Traoré D, Aké-Assi L. Etude botanique des plantes emménagogues utilisées en médecine traditionnelle par les Abbey et Krobou d'Agboville (Côte-d'Ivoire). Rev Med Pharm Afr. 2008;21:43–60.
 20. Zubaida Y, Ying W, Elias B. Phytochemistry and pharmacological studies on *Solanum torvum* Swartz. Jour of Applied Pharma Sci. 2013;3(4):152-160.
 21. Irie-N'guessan AG, Kablan BJ, Kouakou-Siransy NG, Leblais V, Champy P. Evaluation de la toxicité de cinq plantes antiasthmatiques de la médecine traditionnelle ivoirienne. Int. J. Biol. Chem. Sci. 2011;5(3):1316-1319.
 22. Coulerie P. Etude phytochimique et pharmacologique de plantes de Nouvelle-Calédonie à potentialités anti-dengue. Thèse de Doctorat de chimie des substances naturelles. Université de la Nouvelle-Calédonie : Ecole Doctorale du Pacifique. 2012;296.
 23. Busser C. Baies, fruits et pseudo-fruits toxiques utilisés en médecine populaire ou en phytothérapie Phytot. 2007;1:31–36.

© 2018 Yao et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/26685>