



# **Impact of Foliar Application of Boron on Yield and Yield Attributes, Assimilate Supply, Production Efficiency and Fertilizers Productivity of Indian Mustard (*Brassica juncea*) in Field Experiments**

**Samar Pal Singh <sup>a\*</sup>, D. K. Rana <sup>a</sup>, Brijesh Yadav <sup>a</sup>, Kailash <sup>a</sup> and Jai Parkash <sup>a</sup>**

<sup>a</sup> *Krishi Vigyan Kendra, New Delhi, 110073, India.*

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/jeai/2024/v46i72559>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117367>

**Original Research Article**

**Received: 28/03/2024**

**Accepted: 31/05/2024**

**Published: 04/06/2024**

## **ABSTRACT**

Field experiments were conducted for three consecutive *rabi* seasons from 2018 to 2020 to assess the impact of foliar application of boron fertilizer on yield and yield attributes, assimilate supply, production efficiency and fertilizers productivity of Indian mustard [*Brassica juncea* (L.) Czern. & Coss.]. The results showed that the number of primary branches/ plant (7.00 to 8.08), number of

\*Corresponding author: E-mail: [samarpalagro@gmail.com](mailto:samarpalagro@gmail.com);

siliquae/plant (427.2 to 468.8), number of seeds/siliqua (13.03 to 14.01), test weight of seed (3.60 to 3.80 g) and plant dry matter production (38.10 to 41.20 g/plant) were significantly increased with T2 treatment (100% NPKS + foliar application of Boron @ 0.5% Borax) as compared to farmer's practice T1 (100% NPKS + without boron) during three years of experiments. Significantly higher seed yield (22.5 to 24.7 q/ha) of the mustard crop was also reported with T2 (100% NPKS + foliar application of Boron @ 0.5% Borax) as compared to farmers' practice T1 (100% NPKS + without boron) (seed yield 19 to 22.5 q/ha) during the studies. Similarly, Treatment T2 resulted in significantly higher production efficiency and assimilates supply of mustard crop than that of T1 (farmer's practice). The fertilizer productivity in the mustard crop that is directly related to seed yield and the partial factor of productivity of N, P, K and S fertilizers ( $PF_{NPKS}$ ) were also increased significantly with the T2 compared to farmers practice T1 (farmer's practice), during 3 years of studies. The highest gross return and net return and benefit-cost ratio were obtained with T2 (100% NPKS + foliar application of Boron @ 0.5% Borax) treatment over T1 (100% NPKS + without boron).

**Keywords:** Assimilate supply; economics; fertilizers productivity; production efficiency and seed yield.

## 1. INTRODUCTION

Indian mustard (*Brassica juncea*) stands as one of the principal oilseed crops globally, revered for its nutritional value, adaptability, and economic significance. In agricultural landscapes, maximizing the yield and quality of mustard crops is imperative to meet the growing demands of food, feed, and industrial applications. Among the various factors influencing crop productivity, the availability and uptake of essential micronutrients play a pivotal role in determining plant growth, development, and ultimately, yield. Boron is an essential micronutrient for vascular plants and necessary for the growth of crop plants [1]. Low availability of boron in soils impairs the development of floral organs and the elongation of pollen tubes, which leads to a loss of crop yield [2]. Boron deficiency is the second most important micronutrient constraint in crop production after zinc [3]. Rapeseed–mustard is extremely sensitive to boron (B) deficiencies [2]. Boron deficiency in soil leads to the erratic growth of seedlings and reduced photosynthesis [4]. Boron is an important component for various biological processes in plants, including the growth and development of pollen tubes, the maintenance of membrane integrity, pollination and seed production [5]. The primary roles of boron is include the breakdown of nucleic acids, carbohydrates, proteins, indole acetic acid, and phenol, which are involved in the synthesis of plant cell walls and the maintenance of membrane integrity [6]. Additionally, boron plays a key role in cellular division and the control of carbohydrate and protein metabolism, and these processes influence the reproductive phase and development of seeds [7]. Boron is closely associated with plant growth and plays a vital

role in cell division as well as synthesis of oil and protein. Flowering and pods formation is restricted due to boron deficiency and rapeseed-mustard crops are very responsive to boron fertilizer [8,9]. Lu et al. [10] and [11] found that boron fertilizer contributed 611 kg/ha (48.5%) yield advantage of the rapeseed. Yield reduction due to boron deficiency often occurs even if specific visual symptoms of leaves are not apparent. The majority of researchers have focused on foliar boron management in mustard in small experimental plots thus far; consequently, work needs to be done under field conditions to evaluate the influence at the field level. Keeping in mind the problem of low yield of mustard due to boron deficiency in the soil, an on farm trial was designed on foliar application of boron fertilizer in mustard to reduce the yield loss due to the deficiency of boron and to obtain higher yield and quality of mustard crop.

## 2. MATERIALS AND METHODS

### 2.1 Experimental Details


On -farm trials on foliar application of boron on Indian mustard were carried out at the farmer's fields of Qazipur, Shikarpur and Jhatikara villages of NCT Delhi under similar agro climatic conditions during the *rabi* season from 2018 to 2020 (Fig. 6). The rainfall and average temperature of the experimental area from 2018 to 2021 have been depicted in Figs. 3,4,5. The experimental soil was a sandy loam in texture, low in organic carbon (0.31%), available nitrogen (225 Kg/ ha), phosphorous (21 Kg/ ha), potash (195 Kg/ha). The experiments were laid out on five farmer's fields with 5 replications in a 1000 m<sup>2</sup> plot area each and having 2 treatments i.e T1

Farmer’s practice (100% NPKS + without boron) and T2 (100% NPKS + foliar application of Boron @ 0.5% Borax).


### 2.2 Crop Management

The recommended dose of fertilizer was 80:60:40: 30 kg N, P, K,S ha<sup>-1</sup>, respectively, for Indian mustard. Nutrients were supplied through urea, single super phosphate, muriate of potash and bentonite sulphur. Boron was supplied through borax which contains 11% boron through foliar application. Whole of P, K and S and half of the N were supplied as basal dose and the remaining half of the nitrogen was applied 30 days after sowing. Proper weed management, insect- pest management and irrigation management was done in experimental fields.




The Number of primary branches and total siliquae/ plant was counted from five randomly selected plants at harvest and the mean value was taken. The numbers of seeds per siliquae is also counted from randomly selected siliquae for the selected plants. Four plants were selected randomly from the observational rows of each plot and dug out with the help of *khurpi* at 60 DAS, 90 DAS and at harvest. These plants were sun-dried for 48 hours. After sun drying, these plants were dried in the oven at 65±5°C temperature for 48-72 hours or till the samples attained a constant weight, and weighed. The dry matter was expressed in g plant<sup>-1</sup>. The data output was collected from experimental fields as well as farmer’s and cost of cultivation, net income, and benefit -cost ratio were also worked out.



**Fig. 1. Effect of foliar application of boron on cost of cultivation, gross return and net return of the mustard crop during 2018 -2020**



**Fig. 2. Effect of foliar application of boron on benefit- cost ratio of the mustard crop during 2018 -2020**



**Figs. 3,4,5. Rainfall distributions (mm) during crop growing *rabi* season 2018 -21**



**Fig. 6. Site of Krishi Vigyan Kendra in NCT Delhi and its working area**

### 2.3 Crop Efficiency Parameters

Partial factor productivity of nitrogen (PFP<sub>N</sub>), phosphorus (PFP<sub>P</sub>) and Potash (PFP) fertilizer was calculated as per the following formula [12].

$$\text{Partial factor of productivity (PFP}_{\text{npk}}) = \frac{\text{Seed yield (kg)}}{\text{Rate of fertilizer applied (kg)}}$$

Assimilate supply (AS) is the ratio of above ground dry matter of individual plant / number of siliquae. Production efficiency (PE), it represents the increase in seed yield on a daily basis and calculated by dividing total mustard seed production ha<sup>-1</sup> in a sequence with total duration of the crops in a sequence [13].

### 2.4 Statistical Analysis

The yield data were collected from both the experiment and farmers' practice and analyzed by using simple statistical tools. All data were analyzed were using ANOVA, and two sample test values at 5% level of significance were calculated and used to test significance difference between treatment means.

## 3. RESULTS AND DISCUSSION

### 3.1 Yield Attributes

Foliar application of boron had a significant impact on yield attributes of the mustard crop

such as number of primary branches per plant, number of siliquae/plant, and number of seeds/siliqua and 1000-seed weight during 3 years of studies (Table 1). Significantly higher number of primary branches/ plant (7 to 8.08), number of siliquae/plant (427.2 to 468.8), number of seeds/siliqua (13.03 to 14.01) were obtained with T<sub>2</sub> treatment (100% NPKS + foliar application of Boron @ 0.5% Borax) as compared to farmer's practice T<sub>1</sub> (100% NPKS + without boron) during three years of experiments. Similarly, the higher test weight of seed (3.6 to 3.8 g) was reported with the with T<sub>2</sub> treatment (100% NPKS + foliar application of Boron @ 0.5% Borax) in mustard crop, which was significantly higher over farmer's practice T<sub>1</sub> (100% NPKS + without boron) in which test weight of seed was 3.2 to 3.3 g. The partial or complete failure of boron fertilization leads to pollen abortion, seed drop, deformed and shrunken seeds and partially filled pods resulting in lower test weight [14] and [15]. Additionally, Jankowski et al. [16] observed that foliar boron application in mustard crops resulted in considerably larger numbers of primary branches/ plant, siliquae /plant, seeds/ siliqua, and 1000-seed weight. The highest values of yield attributes in treatment T<sub>2</sub> might be attributed to the involvement of boron in hormone synthesis and translocation, carbohydrate metabolism and the synthesis of DNA in plants

[17]. Identical results for plant growth parameters were reported by Shah et al. [18] and Handiganoor et al. [19].

### 3.2 Plant Dry Matter Production

Foliar application of boron had a notable impact on dry matter accumulation during 3 years of the studies (Table 2). The dry matter production (38.1 to 41.2 g/plant) in mustard crop was significantly higher with T<sub>2</sub> treatment (100% NPKS + foliar application of Boron @ 0.5% Borax) as compared to the farmer's practice T<sub>1</sub> (100% NPKS + without boron), in which an increase of 36.4 to 38 g/plant dry matter was recorded. The dry matter production was found to be increased could be attributed to the increased number of branches per plant arising out of the better growth and development conditions facilitated by proper supply of photosynthates to the plants parts. Boron is needed in small quantity by plants for division and cell enlargement in meristem tissue, then the meristem generate new cells at the tip of the root or stem that resulted in increased plant growth or length. The increase growth of plant and number of leaf affects the light received by the plant so that it can improve the performance of photosynthesis and produce many photosynthate in forms of starch, lipids, and proteins. Photosynthate is translocated inside the plant which helps to increase the dry matter of the plant [20].

### 3.3 Seed Yield

The seed yield of the mustard crop was influenced by foliar application of boron during 3 years of studies. The significantly higher seed yield (22.50 to 24.70 q/ha) of the mustard crop was reported with T<sub>2</sub> treatment (100% NPKS + foliar application of Boron @ 0.5% Borax) as compared to the farmer's practice T<sub>1</sub> (100% NPKS + without boron), in which the seed yield was 19.00 to 22.50 q/ha during the experiments (Table 2). These results corroborated well with the studies conducted by Jankowski et al. [16] and Zhao et al. [3] in which they reported higher seed yield with boron in mustard field. Lu et al. [10] and [11] found that boron fertilizer contributed 611 kg/ha (48.5%) yield advantage of the rapeseed. The increase in the seed yield can be ascribed to boron as it is directly linked with the process of fertilization, pollen producing capacity of anther, viability of pollen grains, pollen germination and pollen tube growth [21] and [22]. Boron's involvement in maintaining the

structural integrity of plasma membrane, metabolism of carbohydrates, and synthesis of DNA probably resulted in the additional growth in crops [19]. The reason for the lowest seed yield under farmers practice, might be the higher pollen infertility and lower seed filling as they play an extremely crucial role in both the processes [23]. Correlation analysis showed that seed yield had a significant positive correlation with number of siliquae, seeds/ siliqua, test weight, dry matter accumulation, assimilate supply, however, negative correlation was observed with primary branches (Table 4).

### 3.4 Production Efficiency

The production efficiency of a crop refers to the daily increase in seed yield. Foliar application of boron exerted significant effects on the production efficiency of mustard crop during 3 years of studies (referencing Table 2). Specifically, as a result of foliar application of boron (T<sub>2</sub> treatment), the production efficiency of the mustard crop became significantly higher (16.67 to 18.31) than that of the farmer's practice T<sub>1</sub> (100% NPKS + without boron), whose production potential was found to be 14.07 to 16.67. Foliar application of boron helps in proper translocation of photosynthates toward the seeds and siliquae formation, which enhances in the seeds and siliquae production in mustard. Whereas, the seeds and siliquae development was poor under farmers practices in which boron fertilizer was not applied. The translocation of food material from source to sink was better in boron spray treatment as compared to farmer's practices.

### 3.5 Assimilate Supply

Assimilate supply was influenced significantly with the application of boron fertilizer has been depicted in Table 2. Assimilate supply refers to a plant's ability to mobilize photosynthates once again and it is the ratio of above ground dry matter of individual plant to the number of siliquae formation. During the 3 years studies, assimilate supply of the mustard crop was increased significantly (11 to 11.76) with T<sub>2</sub> treatment (100% NPKS + foliar application of Boron @ 0.5% Borax) over farmer's practice T<sub>1</sub> (100% NPKS + without boron) in which it was increased up from 10.33 to 11.17. At the time of the peak flowering period, the nutrients and carbohydrates begin to remobilize toward seed setting and siliquae formation. Similar finding were also confirmed by Shekhawat et al. [24].

**Table 1. Effect of foliar application of boron on number of primary branches, number of siliquae /plant, number of seeds / siliqua, test weight (g) of the mustard crop during 2018 -2020**

Treatment	Number of primary branches /plant			Number of siliquae / plant			Number of seeds/ siliqua			Test weight(g)		
	2018	2019	2020	2018	2019	2020	2018	2019	2020	2018	2019	2020
T1- Farmer's Practice (100%NPK+ without boron)	5.00±0.8	5.50±1.1	5.5±1.1	402.5±8.2	398.0±8.6	400.8±12.4	11.75±0.47	10.95±0.72	12.01±0.88	3.2±0.23	3.37±0.30	3.25±0.35
T2- (100% NPKS + foliar application of Boron @ 0.5% Borax)	7.75±0.5	7.00±0.7	8.08±0.8	468.8±26.9	427.2±26.6	451.3±36.4	13.53±0.76	13.03±0.65	14.01±0.48	3.6±0.28	3.70±0.63	3.85±0.33
P(T<=t)	0.001	0.020	0.016	0.002	0.022	0.024	0.001	0.005	0.001	0.005	0.027	0.028

**Table 2. Effect of foliar application of boron on dry matter production (g/plant) seed yield (q/ha), assimilate supply and production efficiency of the mustard crop during 2018 -2020**

Treatment	Dry matter production/plant (g)			Seed yield (q/ha)			Assimilate Supply (g siliqua <sup>-1</sup> )			Production efficiency (kg ha <sup>-1</sup> day <sup>-1</sup> )		
	2018	2019	2020	2018	2019	2020	2018	2019	2020	2018	2019	2020
T1- Farmer's Practice (100%NPK+ without boron)	38.4±1.8	36.4±4.6	38.9±1.7	22.5±0.9b	19.0±2.0b	20.0±1.7b	10.51±0.79	11.17±0.61	10.33±1.11	16.67±0.70	14.07±1.49	14.81±1.30
T2- (100% NPKS + foliar application of Boron @ 0.5% Borax)	39.9±1.2	38.1±1.1	41.2±1.4	24.7±0.6a	22.5±0.9a	24.0±0.7a	11.76±1.8	11.22±0.66	11±0.60	18.31±0.51	16.67±0.70	17.78±0.64
P(T<=t)	0.108	0.221	0.044	0.007	0.021	0.001	0.026	0.480	0.203	0.001	0.041	0.001

**Table 3. Effect of foliar application of boron on partial factor of productivity of N,P, K and S of the mustard crop during 2018 -2020**

Treatment	N Partial factor productivity (kg kg <sup>-1</sup> )			P Partial factor productivity (kg kg <sup>-1</sup> )			K Partial factor productivity (kg kg <sup>-1</sup> )			S Partial factor productivity (kg kg <sup>-1</sup> )		
	2018	2019	2020	2018	2019	2020	2018	2019	2020	2018	2019	2020
T1- Farmer's Practice (100%NPK+ without boron)	28.13±1.19	23.75±2.5	25±2.19	37.50±1.5	31.67±3.3	33.3±2.92	56.25±2.39	47.50±5.5	50±4.38	75.00±3.19	63.33±6.73	66.67±5.85
T2- (100% NPKS + foliar application of Boron @ 0.5% Borax)	30.90±0.87	28.13±1.1	30±1.08	41.19±1.1	37.50±1.4	40.0±1.44	61.79±1.74	56.25±2.40	60±2.16	82.39±2.32	75.00±2.28	80±2.88
P(T<=t)	0.007	0.021	0.001	0.027	0.001	0.005	0.007	0.021	0.001	0.002	0.011	0.005

**Table 4. Relationship between yield and yield attributes of mustard as influenced by boron spary (mean data of three years)**

X-axis	Y-axis	Correlation coefficient R <sup>2</sup>	Regression equation
Grain yield	Primary Branches	0.57	y = 0.447x - 3.4144
Grain yield	Number of siliquae/ plant	0.77	y = 11.739x + 165.11
Grain yield	Number of seeds/ siliqua	0.75	y = 0.4563x + 2.4576
Grain yield	Test weight (1000 seed weight)	0.39	y = 0.0736x + 1.8658
Grain yield	Drymatter accmualtion	0.60	y = 0.5695x + 26.223
Grain yield	Assimilate supply	0.22	y = 0.1108x + 8.5482



Application of boron might be involved in the synthesis of protein, chloroplast pigments, and electron transfer system that led to the increased photosynthetic activity as well as assimilate supply of mustard crop [25]. The positive effect of boron application has been reported on assimilate supply by Haque et al. [26] and Islam [8].

### 3.6 Fertilizer Productivity

Foliar spray of boron had a significant impact on NPK & S fertilizers productivity during 3 years of studies. Fertilizer productivity was evaluated in terms of partial factor of productivity (PFP) and calculated based on the yield of the mustard crop (Table 3). Over the course of study years, the PFP of N, P, K and S fertilizers (PFP<sub>NPKS</sub>) were increased significantly (28.13 to 30.9, 37.5 to 41.19, 56.25 to 61.79, 75 to 82.39 kg kg<sup>-1</sup>) with T<sub>2</sub> treatment (100% NPKS + foliar application of Boron @ 0.5% Borax) as compared to the farmer's practice T<sub>1</sub> (100% NPKS + No use of boron). It was observed that the PFP of S was found to be higher followed by PFP of K over of PFP of N and P. It means boron had a positive interaction with other nutrients like N, P, K and S, and plays an important role to increase the fertilizer productivity of NPK&S. Higher PFP of NPK&S with a foliar application of boron might be ascribed for higher seed yield production and better translocation of food materials toward seeds development than without boron application field. All these are in accordance with the findings reported by Singh et al. [12] and [2].

### 3.7 Economics

The economics of the treatments was calculated depending on the prevailing market prices of the inputs and outputs for the particular year. During 3 years of studies, the higher gross return of Rs. 85500 to 111600/ha, net return of Rs.63210 to 72828/ha and B: C ratio 2.83 to 4 were obtained with T<sub>2</sub> treatment (100% NPKS + foliar application of Boron @ 0.5% Borax) as compared to farmer's practice T<sub>1</sub> (100% NPKS + without boron) has been illustrated in Fig. 1. The higher benefit- cost ratio with T<sub>2</sub> treatment was found due to the higher yield obtained under T<sub>2</sub> treatment as compared to T<sub>1</sub> (farmer's practice) (Fig. 2). Hence, higher benefit- cost ratio proved the economic viability of the technology intervention and convinced the farmers about the utility of improved technologies. Similar economic benefits owing to adoption of foliar application of

boron in mustard were also reported by Dubey et al. [27].

## 4. CONCLUSION

Based on the results of on-farm trials conducted in real farming conditions, it can be concluded that the foliar application of boron, specifically at a concentration of 0.5% borax alongside 100% recommended NPKS fertilizers, enhances the yield attributes of the mustard crop. Additionally, the foliar application of boron has been shown to improve the production efficiency and assimilate supply of the mustard crop. Moreover, the fertilizer productivity of N, P, K, and S fertilizers was observed to increase with the foliar application of boron at the mentioned concentration over the course of three years of studies. Consequently, foliar application of boron at 0.5% borax concentration was identified as optimal for achieving higher seed yields in boron-deficient soils. The adoption of foliar application of boron has the potential to significantly augment both the income and livelihoods of farming communities.

## ACKNOWLEDGEMENTS

The financial support to meet the expenses towards frontline demonstrations by the Department of Agricultural & Farmers Welfare, Government of India through its nodal agency ICAR-Agricultural Technology Application Research Institute (ATARI), Jodhpur, Rajasthan, India is gratefully acknowledged [28].

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Wimmer MA, Abreu I, Bell RW, Bienert MD, Brown PH, Dell B, Fujiwara T, Goldbach HE, Lehto T, Mock HP, Wirén N. Boron: an essential element for vascular plants. *New Phytologist*. 2019;226(5): 1232-7.
2. Zhao Z, Wang Y, Shi J, Wang S, White PJ, Shi L, Xu F. Effect of balanced application of boron and phosphorus fertilizers on soil bacterial community, seed yield and phosphorus use efficiency of Brassica napus. *Science of the total environment*. 2021;10,751:141644.

3. Rehman A, Wang X, Hussain A, Qamar R, Khalofah A, Hussain M. Boron Application in Yermosols Improves Grain Yield and Quality of Chickpea (*Cicer arietinum* L.). Journal of King Saud University-Science. 2022;1,34(2):101768.
4. Safdar ME, Qamar R, Javed A, Nadeem MA, Javeed HM, Farooq S, Głowacka A, Michałek S, Alwahibi MS, Elshikh MS, Ahmed MA. Combined application of boron and zinc improves seed and oil yields and oil quality of oilseed rape (*Brassica napus* L.). Agronomy. 2023;29,13(8):2020.
5. Ahmad W, Zia MH, Malhi SS, Niaz A, Ullah S. Boron Deficiency in soils and crops: a review. Crop plant. 2012;20;2012:65-97.
6. Goldbach HE, Yu Q, Wingender R, Schulz M, Wimmer M, Findeklee P, Baluška F. Rapid response reactions of roots to boron deprivation. Journal of Plant Nutrition and Soil Science. 2001;164(2):173-81.
7. Miwa K, Fujiwara T. Boron transport in plants: co-ordinated regulation of transporters. Annals of Botany. 2010;1, 105(7):1103-8.
8. Islam MB. Requirement of boron for mustard, wheat and chickpea based rice cropping patterns. PhD diss., Department Soil Science, Bangladesh Agricultural University, Mymensingh, Bangladesh. 2005;1-124
9. Saha PK, Saleque MA, Zaman SK, Bhuiyan NJ. Response of mustard to S, Zn and B in calcareous soil. Bangladesh J. Agril. Res. 2003;28(4):633-6.
10. Lu X, Lu Y, Mao D. The levels of available boron in red earth in central Zhejiang Province and boron nutrition of rape. Soils and Fertilizers (Beijing). 2000;(1):30-4.
11. Randhawa MK, Dhaliwal SS, Sharma V, Toor AS, Sharma S, Kaur M, Verma G. Nutrient use efficiency as a strong indicator of nutritional security and builders of soil nutrient status through integrated nutrient management technology in a rice-wheat system in northwestern India. Sustainability. 2021;20,13(8):4551.
12. Singh SP, Mahapatra BS, Pramanick B, Yadav VR. Effect of irrigation levels, planting methods and mulching on nutrient uptake, yield, quality, water and fertilizer productivity of field mustard (*Brassica rapa* L.) under sandy loam soil. Agricultural Water Management. 2021;1,244:106539.
13. Premi OP, Kandpal BK, Rathore SS, Shekhawat K, Chauhan JS. Green manuring, mustard residue recycling and fertilizer application affects productivity and sustainability of Indian mustard (*Brassica juncea* L.) in Indian semi-arid tropics. Industrial Crops and Products. 2013; 1(41):423-9.
14. Kapila Shekhawat KS, Shivay YS, Dinesh Kumar DK. Productivity and nutrient uptake of spring sunflower (*Helianthus annuus*) as influenced by nitrogen sources, sulphur and boron levels, Indian Journal of Agricultural Science. 2008;78:90-94
15. Kumararaja P, Premi OP, Kandpal BK. Application of boron enhances Indian mustard (*Brassica juncea*) productivity and quality under boron deficient calcareous soil in semi-arid environment. 2015;S249-S254.
16. Jankowski KJ, Załuski D, Sokólski M. Canola-quality white mustard: Agronomic management and seed yield. Industrial Crops and Products. 2020;1;145:112138.
17. Verma A, Shahi UP, Chaurasia P, Kumawat L. Effect of mode of micronutrients application on growth and yield of pigeon pea (*Cajanus cajan* L.) in sandy loam soil. Int. J. Curr. Microbiol. App. Sci. 2020;9(03):17-24.
18. Shah KA, Gurjar R, Parmar HC, Sonani VV. Effect of sulphur and zinc fertilization on yield and quality of pigeonpea in sandy loam soil. Green Farming. 2016;7(2):495-7.
19. Handiganoor M, Patil S, Vasudevan S. Response of pigeonpea (*Cajanus cajan* L.) to seed polymerization with micronutrients and foliar spray at different growth stages. British Journal of Environment and Climate Change. 2017;16,7(4):205-13.
20. Timotiwu PB, Agustiansyah A, Pramono E, Agustin S. Phosphorus and boron application on growth, yield, and quality of soybean seeds (*Glycine max* [L.] Merril). Journal of Agricultural Science. 2018;1,10(6):372-82.
21. Padbhushan R, Kumar D. Yield and nutrient uptake of green gram (*Vigna radiata* L.) as influenced by boron application in boron-deficient calcareous soils of Punjab. Communications in soil science and plant analysis. 2015;12, 46(7):908-23.
22. Jaiswal AD, Singh SK, Singh YK, Singh S, Yadav SN. Effect of sulphur and boron on yield and quality of mustard (*Brassica juncea* L.) grown on Vindhyan red soil. Journal of the Indian Society of Soil Science. 2015;63(3):362-4.

23. Hussain M, Khan MA, Khan MB, Farooq M, Farooq S. Boron application improves growth, yield and net economic return of rice. *Rice Science*. 2012, 1;19(3):259-62.
24. Shekhawat K, Rathore SS, Kandpal BK, Premi OP, Singh D, Chauhan BS. Crop establishment techniques affect productivity, sustainability, and soil health under mustard-based cropping systems of Indian semi-arid regions. *Soil and Tillage Research*. 2016;1(158):137-46.
25. Srinivasan G, Gobi R, Balasubramanian A, Sathiyamurthi S. Influence of nipping and nutrient management practices on growth, yield attributes and yield in Pigeonpea. *Plant Archives* .2019;19:737–740.
26. Haque MA, Jahiruddin M, Islam MR. Effect of sulphur and boron on seed yield of mustard (*Brassica napus*). *Bangladesh J. Seed Sci. & Tech*. 2000;4(1&2):7-11.
27. Dubey SK, Gautam US, Singh AK, Singh A, Chahal VP, Singh AK, Singh CH, Srivastava AJ. Quantifying the yield gap minimization in lentil (*Lens culinaris*) under Cluster Frontline Demonstrations (CFLD) conducted in Uttar Pradesh. *Indian Journal of Agricultural Sciences*. 2018;1;88(6):851-9.
28. GOI (Government of India). *Agricultural Statistics at a Glance*. Agricultural Statistics Division, Department of Agriculture and Cooperation and Farmers Welfare, Ministry of Agriculture, GOI, New Delhi; 2022.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/117367>