



Effect of Neem Cake, Neem Oil and Plant Extracts on Purple Blotch of Onion (*Allium cepa* L.) Caused by *Alternaria porri* (Ellis) Cif.

Arunesh Singh ^{a*} and Abhilasha A. Lal ^a

^a Department of Plant Pathology, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, (Uttar Pradesh), India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i102984>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124655>

Original Research Article

Received: 06/08/2024

Accepted: 08/10/2024

Published: 22/10/2024

ABSTRACT

The present study was conducted to evaluate the Effect of neem cake, plant extracts and neem oil on purple blotch of onion (*Allium cepa* L.) caused by *Alternaria porri* (Ellis) Cif. Six different treatments were tested in a randomized block design with 3 replications at the Central Research Field, Department of Plant Pathology, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj during Rabi season of 2023. The results should that the minimum per cent disease intensity was recorded in T₁Neem cake + Neem oil (31.72), T₅ Neem cake + Eucalyptus leaf extract (37.40). bulb weight of onion (71.13g), bulb diameter (7.13cm), bulb yield per plot (2.106kg), yield (105.317 q/ha) and benefit cost ratio (1: 4.42) were found in T₁Neem cake + Neem oil (31.72) when compared to untreated check and other treatments.

*Corresponding author: E-mail: aruneshsingh1244@gmail.com;

Cite as: Singh, Arunesh, and Abhilasha A. Lal. 2024. "Effect of Neem Cake, Neem Oil and Plant Extracts on Purple Blotch of Onion (*Allium cepa* L.) Caused by *Alternaria porri* (Ellis) Cif.". *Journal of Experimental Agriculture International* 46 (10):612-17. <https://doi.org/10.9734/jeai/2024/v46i102984>.

Keywords: *Alternaria porri*; neem cake; neem oil; onion; plant extract; purple blotch.

1. INTRODUCTION

Onion (*Allium cepa* L.) is one of the most important vegetable crops belongs to the family Amaryllidaceae. There are more than 780 species, usually perennial bulbous plants grown worldwide [1].

Onion is also called as "Queen of kitchen". It contains a phytochemical called quercetin which is effective in reducing the risk of cardiovascular diseases, anticancer and has antioxidant properties. It is supposed to have its origin in the middle East Asian countries and introduced in India from Palestine [2].

Production of Onion in 2022 is expected to be around 254.73 lakh tone (APEDA, 2022). India the yield of onion is very low as compared to the world average yield of 19.1t/ha [3].

The most important factors responsible for the diseases like purple blotch, downy mildew, stem phylum blight, basal rot and storage rots. Among the foliar diseases, purple blotch is one of the most destructive diseases, commonly prevailing in almost all onion growing pockets of the world, which causes heavy loss in onions under field conditions. Losses can range from 30 to 100 per cent. The disease may reach epidemic states during the favorable conditions of high relative humidity (80-90%) and optimum temperature ($24\pm 10^{\circ}\text{C}$). Purple blotch disease of onion causes significant reduction in foliar production and bulb yield [4].

The several effective fungicides have been recommended against this pathogen, but they

are considered as a long-term solution, due to concerns of expense, exposure risks and the hazards of its residues. Moreover, the development of resistance of pathogenic fungi towards synthetic pesticides is of great concern that can affect significantly the efficacy of chemical fungicides. The present study was aimed to determine the efficacies of different bio-agents and neem oil against purple blotch of onion [5].

2. MATERIALS AND METHODS

The study was conducted in randomized block design (RBD) with plot size $2\times 1\text{m}^2$ and there were three replications for each treatment. The experiment was conducted on Nasik red variety of onion. Observations on disease intensity of purple blotch of onion were recorded at 30 days interval, bulb weight of onion (g), bulb diameter (cm), bulb yield per plot (kg), yield (q/ha) and B:C ratio data were obtained after the harvest on physiological maturity.

2.1 Preparation of Plant Extracts

Hundred grams of fresh healthy plant parts (leaves) collected from field was washed with distilled water and air dried and crushed in 100 ml of distilled water (w/v). The crushed product was filtered through double layer, muslin cloth and further filtrated through Whatman No. 1 filter paper using funnel and volumetric flasks (100 ml cap.). The prepared solution was 100 per cent concentration, which was further diluted to required concentrations of 10 per cent [6].

Table 1. Treatments combination

Treatment Number	Treatment Details	References
T ₀	Control (Untreated)	-
T ₁	Neem cake @440gm (S.A) + Neem oil 5% (F.S)	Kumar (2007)
T ₂	Neem cake @440g (S.A) + Datura leaf extract @10% (F.S)	Brahmane et al. (2015)
T ₃	Neem cake @440g (S.A) + Lantana leaf extract @10% (F.S)	Brahmane et al. (2015)
T ₄	Neem cake @440g (S.A) + Bael leaf extract @10% (F.S)	Kumar (2007)
T ₅	Neem cake @440g (S.A) + Eucalyptus leaf extract@10% (F.S)	Kumar (2007)
T ₆	Mancozeb @0.2% (F.S) (Treated check)	Chethana et al. [5]

Dose: Neem cake @440g 2m²(S.A), Plant extract (10%), neem oil (5%) and mancozeb (0.2%)
S.A: - Soil amendment; F.S: - Foliar spray

2.2 Percent Disease Intensity (PDI)

Percent disease index was calculated by using the following formula [7].

$$\text{Percent Disease Index (PDI)} = \frac{\text{Sum of individual ratings}}{\text{No of leaves observed}} \times \frac{100}{\text{Maximum disease grade}}$$

Table 2. Disease rating scale

Scale	Description
0	No disease symptoms
1	A few spots towards tip covering 10% leaf area
2	Several dark purplish brown patches covering up to 20% leaf area.
3	Several patches with paler outer zone covering up to 40% leaf area.
4	Leaf streaks covering up to 75% leaf area or breaking of the leaves from center.
5	Complete drying of the leaves or breaking of the leaves from center.




Fig. 1. Disease rating scale

To calculate the PDI, five plants were randomly examined from each plot and scored for disease severity using the 0-5 scale established by Sharma, [8].

3. RESULTS

3.1 Effect of Neem Cake, Plant Extracts and Neem Oil on Disease Intensity of Purple Blotch of Onion

The stastically analyzed data presented in Table 3 and depicted in reveald effect of organic amendment, plant extracts and neem oil on disease intensity of *Alternaria porri* on

onion under field condition. The treatments were found to have significantly reduced disease intensity as compared to the control (untreated ckeck). As the treated check (T₆) mancozeb @0.2% reduced the disease intensity of purple blotch of onion at its maximum point within the treatments Neem cake + Neem oil (T₁) showed maximum reduction of disease intensity of Neem cake + Eucalyptus leaf extract (T₅), Neem cake +Lantana leaf extract (T₃),Neem cake + Bael leaf extract (T₄)and Neem cake + Datura leaf extract (T₂)the highest disease was recorded in cntrol (T₀) (Untreated check) with (59.19) % disease intensity.

3.2 Effect of Neem Cake, Plant Extracts and Neem Oil on Bulb Weight of Onion (g), Bulb Diameter (cm), Bulb Yield Per Plot (kg) and Yield (q/ha) Of Onion

The data presented in Table 4 and depicted in reveals the response of selected treatments of organic amendment, plant extracts and neem oil on bulb weight of onion(g), bulb diameter (cm), bulb yield per plot (kg) and yield (q/ha) of onion under field conditions. The results indicate that the untreated check T₀ (control) show significantly reduced for all the replications. The maximum data was recorded T₁Neem cake + Neem oil on fresh weight of bulb (68.80gm), diameter of bulb (6.80cm), bulb yield per plot (1.968kg), bulb yield (98.383q/ha) the lowest data was recorded in control(T₀) (Untreated check).

4. DISCUSSION

The probable reasons for such findings may be due to the soil application of neem cake that could have improved the availability of nutrients to the crop by enhancing the mineralization and supply of readily available nutrients to the soil microbial community which may have supported the plant growth by Sharma et al. [9]. Whereas, neem oil was used as foliar spray that may have inhibited the emergence of fungal pathogen due to existence of major components such as azadirachtin, saladin, nimbi, nimonol and iso-meldenin [10] which have the ability to disrupt the cell membrane and facilitate intracellular compound leakages lead to cell lysis resulting in healthy plant growth. Similar findings have been reported by Sindu et al. [11] and Divya et al. [12]

Table 3. Effect of neem cake, neem oil and plant extracts on disease intensity of purple blotch of onion

Treatment Number	Treatment Details	Disease intensity (%)		
		30 DAT	60 DAT	90 DAT
T ₀	Control (Untreated)	10.41	38.60	59.19
T ₁	Neem cake + Neem oil	8.11	19.43	31.72
T ₂	Neem cake + Datura leaf extract	9.45	27.40	45.31
T ₃	Neem cake + Lantana leaf extract	8.85	23.44	41.57
T ₄	Neem cake + Bael leaf extract	9.15	25.46	43.60
T ₅	Neem cake +Eucalyptus leaf extract	8.75	21.54	37.40
T ₆	Mancozeb (Treated check)	7.98	15.49	19.46
	CD (5%)	0.90	0.63	0.26

Table 4. Effect of neem cake, plant extracts and neem oil on bulb weight of onion (g), bulb diameter (cm), bulb yield per plot (kg) and yield (q/ha) of onion

Treatment Number	Treatment Details	Bulb weight(g)	Bulb diameter(cm)	bulb yieldper plot (kg)	yield (q/ha)
T ₀	Control (Untreated)	40.43	3.40	1.370	39.817
T ₁	Neem cake + Neem oil	68.80	6.80	1.968	98.383
T ₂	Neem cake + Datura leaf extract	57.86	4.86	1.538	80.900
T ₃	Neem cake + Lantana leaf extract	61.40	5.92	1.751	89.567
T ₄	Neem cake + Bael leaf extract	59.53	5.43	1.645	86.233
T ₅	Neem cake +Eucalyptus leaf extract	64.20	6.15	1.869	93.450
T ₆	Mancozeb (Treated check)	71.13	7.13	2.106	105.317
	CD (5%)	0.16	0.05	0.07	0.83

who revealed that the maximum disease inhibition was found in the neem oil treatment, suggesting that the use of garlic extracts could be a viable approach to control *Alternaria porri*. Though among all the treatments chemical fungicide (Treated check) has shown the minimum disease control with strong fungicidal effect against the pathogen. As of, it may produce some toxic chemical residues, which may have potential harmful effects to non-targeted organisms. So, considering the ecosystem, neem oil significantly inhibits the pathogen could lead to better health of the plants which in turn helps in producing minimum disease intensity and leading to a healthier plant and environment friendly.

5. CONCLUSION

From the results of this study it may be concluded that neem cake (Soil amendment) + neem oil (Foliar spray) recorded the minimum disease intensity of purple blotch of Onion caused by *Alternaria porri* (Ellis) Cif. and bulb weight of onion (g), bulb diameter (cm), bulb yield per plot (kg), yield (q/ha) and benefit cost ratio of onion. The present investigation was limited to one crop season (Rabi, 2023-24), under the agroclimatic conditions of Prayagraj (U.P.) hence, additional research is necessary to validate the findings of the study as the data collected for one crop season is inadequate to make definite conclusions.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mukul Lal AA, Simon S. Effect of oil cakes and lemongrass essential oil on purple blotch (*Alternaria porri*) disease of onion (*Allium cepa* L.). Annals of Plant Protection Sciences. 2024;32(1): 48-53.
2. Kareem M, Abdul Bhat, Amruta S, Kurubetta Krishna D, Mesta RK, Allolli TB, Ajjappalavar PS, Shweta K, Masuthi S, Dileepkumar, Waseem MA. Evaluation of silicon and bio-formulations for the management of purple blotch disease of onion. Research Journal of Chemical and Environment Sciences. 2018;6(1): 46- 50.
3. Bhoite V, Backiyavathy MR. Effect of sources and levels of sulphur on quality of onion (*Allium cepa* L.). International Journal of Current Microbiology Applied Science. 2022; 11(07):206-211.
4. Gupta RBL, Pathak VN. Yield losses in onion due to purple leaf blotch disease caused by *Alternaria porri*. *Phytophylactica*. 1988;20:21-23.
5. Chethana BS, Ganeshan G, Archana SR, Bellishree K. Integrated disease management of purple blotch disease of onion. International Journal of Quarantine Life Sciences. 2015;10(4): 1729-1733.
6. Shekhawat PS, Prasada RL. Antifungal activities of some plant extracts. Inhibition of spore germination. Indian Phytopathology. 1971;24:800-802.
7. Wheeler BEJ. An introduction to plant diseases. John Wiley and Sons Limited, London. 1969;374.
8. Sharma SR. Effect of fungicidal sprays on purple blotch and bulb yield of onion. Indian Phytopathology. 1986;39: 78-82.
9. Sharma SK, Tomar DS, Nayak MK. Screening of fungicides, essential oil and plant extracts against early blight of potato caused by *Alternaria solani*. International Journal of Chemical Studies. 2019;7(6):1166-1170.
10. Hinduja N, Simon S, Lal AA. Effect of selected bio-resources on purple blotch disease of onion (*Allium cepa* L.). The Pharma Innovation Journal. 2021;10(10): 366-370.
11. Sindu PL, Tiwari S, Govardhani P, Supriya V. Efficacy of bio-agents and neem oil against *Alternaria porri* (Ellis)Cif. in onion (*Allium cepa* L.). International Journal of Plant and Soil Science. 2022;34(21): 586-593.

12. Divya Tiwari S, Tiwari JK, Kumar A. Efficacy of different essential oil and bio-agent against purple blotch of onion (*Alternaria porii*). Bangladesh Journal of Botany. 2022;51(3): 541-546.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/124655>