



Holocene Lacustrine Environment in the Western Eritrea-evidence from Freshwater Shells

Wegahta Tesfalidet Berhe^{1*} and Hiroaki Ishiga¹

¹Department of Earth Science, Shimane University, 1060 Nishikawatsu, Matsue, Shimane, Japan.

Authors' contributions

This work was carried out in collaboration between both authors. Author WTB designed the study, performed statistical analysis, literature searched, wrote the protocol and wrote the first draft of the manuscript. Author HI performed sampling, designed and wrote the manuscript with author WTB, managed the laboratory and XRF analysis of the study. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JGEESI/2019/v21i330126

Editor(s):

(1) Dr. Kaveh Ostad-Ali-Askari, Department of Civil Engineering, Isfahan (Khorasgan) Branch, Islamic Azad University, Iran.

Reviewers:

(1) Sciuto Francesco, Catania University, Italy.

(2) Rosario García Giménez, UAM, Spain.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/49443>

Original Research Article

Received 15 March 2019

Accepted 30 May 2019

Published 04 June 2019

ABSTRACT

This study focused on finding of freshwater molluscs in Holocene sediments outcropping in western Eritrea. Three species of gastropods were found in calcareous sediments at the bank of a tributary river named Shatera River, within the license of Bisha Mines, they are *Melanoides tuberculata* (O. F. Müller, 1774), *Bulinus globosus* (Morelet, 1866) and *Indoplanorbis exustus* (Deshayes, 1833). The ¹⁴C dating on the shells of *I. exustus* and *B. globosus* was also carried out. The age dating result showed (9326-9134 cal yr BP) and (9134-8999 cal yr BP) respectively, of which are in the mid-Holocene epoch (11,000–5000 yr BP). This epoch is best known for it was a time when the northern and eastern part of Africa experienced a warm and humid climate. Among the freshwater molluscs, the existence of *Melanoides tuberculata* makes it more significant, for its widespread occurrence in Quaternary deposits throughout Africa and Asia in both fresh and highly evaporated lakes. This is suggestive of the considerable development of the lacustrine environment in the western part of Eritrea, which at present has disappeared.

*Corresponding author: E-mail: wegihtb@gmail.com;

Keywords: *Holocene; freshwater molluscs; lacustrine environment; paleontology; ¹⁴C dating; climate change.*

1. INTRODUCTION

Fossil evidence is important for the determination of age, sedimentary environment and understanding of paleoclimate. The occurrence of fossils of freshwater molluscs (snails) is direct evidence of a lacustrine environment [1,2] and the development of such basin during geologic time. Exposure of bank sediments along some of the tributary rivers in western Eritrea where fluvial and lacustrine sedimentary sequences is developed. Along the banks of one of these rivers, calcareous muds yielded freshwater molluscs. Reports on the freshwater molluscs are rare in northeastern Africa and this discovery is useful for the reconstruction of paleoenvironmental changes [3]. Freshwater molluscs are considered highly sensitive to hydroclimate fluctuations and can record abiotic factors such as water chemistry, depth and turbulence [4,5,6,7].

2. MATERIALS AND METHODS

2.1 Study Area

2.1.1 Geography and geology

Eritrea is situated in the northeastern Africa (Fig. 1). The topography of this country varies in altitude ranging from 60 to more than 3,000 meters above sea level. Eritrea has a variety of climate conditions that range from hot and arid near the Red Sea to sub-humid in isolated micro-catchments along the eastern escarpment, and the central highlands with a pleasant climate. The western lowlands have a comparable climate with the coastal area and December is the coldest month [8]. The mean annual rainfall in the coastal areas is less than 300 mm per year, while in the highlands and the western lowlands, it ranges between 500 and 1,000 mm [9]. The Barka River is one of the major rivers which runs from the highlands to the eastern coast of Sudan (Fig. 2). Several streams flow seasonally eastward from the plateau to reach the sea on the Eritrean coast [9].

Geology of Eritrea comprises of Precambrian basement rocks which cover about 60% of the surface and are unconformably overlain by predominantly Mesozoic sedimentary rocks, and Tertiary to Quaternary volcanic-sedimentary rocks [9]. The major part of the Precambrian rocks consists of Neoproterozoic (ca. 870-670

Ma) continental-marginal and juvenile intraoceanic magmatic-arc rocks. The Neoproterozoic basement rock has been divided into four terranes based on their distinct stratigraphic and structural characteristics [10], namely Barka Terrane, Hagar Terrane, Nakfa Terrane and Arag Terrane (Fig. 1). While the southeastern part of the country, the Danakil segment, and the northeastern part of the country, consists of Cenozoic volcanic and sedimentary rocks [10].

The western segment of the Barka Terrane is developed in the north-western part of the country underlies the Barka lowlands (Fig. 1). It comprises amphibolite, amphibolite-facies, pelitic schists containing kyanite and staurolite, quartzites and marble [10,11].

The central segment, the Hagar Terrane (Fig. 1), contains dominant mafic igneous rocks with various types of meta-igneous and meta-sedimentary rocks [12]. Occasionally, layered chloritic schists occur with epidotic and chloritic meta-basalts. The Hagar Terrane displays an east-verging thrust contact with the adjacent segment to the east. Chromite, platinum group elements, nickel, gold, and copper mineralization are the prospective minerals in this terrane [9].

The eastern segment, Nakfa Terrane (Fig. 1), is made up of granitoid-greenstone belts and syn- to post-tectonic granitoid rocks, varying from gabbro to syeno-diorite to granite [11]. The western Nakfa Terrane, near Bisha Mine (Fig. 1), comprises basin and range style topography with intrusive rocks and a flat valley floor underlined by volcanic and sedimentary rocks [10]. Rock exposures in the valley floor are generally limited to the bases of the adjacent hills [10]. The Nakfa Terrane is the largest of the four Terranes, and it is considered to be a relict of island arc assemblage. Several volcanogenic massive sulfides (VMS), base-metal deposits and gold ores occur in this tectonic unit [10].

The Arag terrane (Fig. 1) is bounded to the east by several Late Tertiary rocks, which consists of granitoid and metasedimentary rocks [12,13].

The southern segment, the Danakil Terrane (Fig. 1), is composed of metamorphic rocks, which may be grouped into three formations; hornblende biotite gneissic formation, a phyllitic formation consisting of schists, conglomeratic

phyllites, crystalline limestones, and graphitic schists; and the third formation is Post-tectonic granitoids [9,10].

As for Quaternary sediments, a thick evaporitic formation with shell material fills the basin in the Danakil Depression. Basaltic flows and related spatter cones represent Quaternary volcanic activity in the Danakil region [11]. However, detail description of the Holocene sediments has not been indicated.

3. RESULTS

3.1 Stratigraphy of the Studied Area

The sampling area is characterized by a stratigraphic sequence consisting of fine-grained sand and mud. Freshwater mollusk occurs in calcareous mud intercalating fine-grained sand overlying conglomerate (Fig. 4), at the riverbank of Shatera River (Fig. 3). This conglomerate consists of angular pebble and cobble within the poorly sorted coarse sandstone. Conglomerate clasts are mainly basic rocks and quartz grains. Shell-bearing bed show light color yellowish grey (10 YR 6/2 of Münsell soil color chart). The bed forms a lenticular body of which thickness decreased laterally rightward (from north to south) from 1 m to 20 cm (Fig. 4). This shell bearing bed changes to poorly sorted sandy silt laterally, which is overlain by massive sandy silt and further by alternating beds of fine-grained sand and mud. Fine-grained sand is characterized by parallel lamination. Three species of gastropods and a species of bivalve have been found. Gastropods shells can be mainly distinguished into two types, elongated conical shell and ovoidal shell which are described below. Elongated and ovoidal shells sporadically occur in the bed, and conical shells occur parallel to the stratigraphic plane (Fig. 5). Ovoidal shells are arranged in the bed of which some are with their apex obliquely upward (Fig. 5). Disk-shaped shells were not common among the observed shells on the mud surface as indicated by the arrow in Fig. 5.

3.2 Sample Collection and Preparation

3.2.1 Shell collection

Samples were collected from three points; A, B and C (Fig. 4) from Shatera Riverbank (Fig. 3), (N 15°31'44.21" and 37°31'24.16"). Besides this, mud of about 263 g was collected from sample point A. These samples were packed in a plastic bag and put in a plastic box, then imported to

Shimane University, Japan for analysis. These samples were imported under the permit of the Ministry of Agriculture, Forestry and Fisheries in accordance with the Plant Protection Law, Japan. Mud sample of about 200 g was softened by water to observe the abundance of the shells within the mud. Relatively larger shells were collected in the field while tiny shells in the laboratory after softening sediments by water. From the collected 3 samples, one-point sample (A1 at SW 6 in Fig. 3) was suitable for the determination of species because of better preservation. Species name and numbers are indicated in Table 1. Smaller shells dominated Table 2 (A1).

3.2.2 Calcareous mud sample collection and procedure for geochemical analysis

Calcareous mud samples A1, A2, and A3 were collected from sample point A (Fig. 4), from Shatera Riverbank. About 200 g calcareous mud samples were packed in plastic bags and imported to Shimane University, Japan for analysis, under the permit of the Ministry of Agriculture, Forestry and Fisheries. Approximately 50 g of each mud sample was dried in an oven for 48 hrs at 160°C to remove weakly-bound volatiles. The dried samples were then grounded for 20 min using automatic agate pestle and mortar grinder. Then for XRF analyses about 5 g of the powdered samples were compressed into briquettes by a force of 200 kN for 60 sec in line with the method of [14]. Average errors for all elements are less than ±10% relatively. Analytical results for GSJ standard JSI-1 were acceptable compared to the proposed values of [15].

3.3 Systematic Paleontology

A. *Melanoides tuberculata* (O. F. Müller, 1774)

Phylum: Mollusca

Class: Gastropoda

Order: Caenogastropoda

Family: Thyridae

Genus: *Melanoides*

Type species: *Melanoides fasciolata* Olivier, 1804 accepted as *Melanoides tuberculata* (O. F. Müller, 1774).

Synonym: *Nerita tuberculata* (O. F. Müller, 1774).

Size: 20–50 mm high and 7-16 mm width [16].

Ecology: *Melanoides tuberculata* is a freshwater benthic species living in and on the bottom sediments of permanent waters, from small springs to vast lakes. Living *Melanoides tuberculata* is known to burrow through lake deposits during the daytime and come out to feed in the night. They are carnivorous and avoid strong current by inhabiting near the banks of freshwater [17].

Geographic distribution: Tropical and sub-tropical regions of Northern Africa and Southern Asia.

Description: *Melanoides tuberculata* (O. F. Müller, 1774) has a dextral turreted (5-17 in mm long) and is with a pointed apex (2-5 in mm wide) of the specimen SW 6-A1 (Fig. 6). The aperture is almost oval and broadens as it goes down to the basal margin of the shell. This species has five to six angular whorls. The inner lip is inconspicuous due to filling by sediments. The sculpture of the shell consists of a parallel row of tubercles of various sizes forming fine transverse lines. Generally, the *Melanoides tuberculata* is of various sizes but have the same shell morphology of the proportion. The occurrence of *Melanoides tuberculata* in the surface of the calcareous mud is higher compared to other species.




Fig. 1. Location of the study area with geologic scheme of Eritrea showing Terrane division [10]




Fig. 2. Map showing Barka River and the location of the study area. Generated using QGIS, 2018




Fig. 3. Satellite image of the study area within Bisha Mine (Google Earth, 2018)




Fig. 4. The outcrop of freshwater molluscs bearing bed (Loc. SW 6 in Figs. 2, 3) within Bisha mining area, Eritrea. A, B and C are indicating shell sample collections of which species and numbers are given in Table 1. A1 in A is for detail shell examination and A1, A2 and A3 are calcareous mud samples collected for geochemical analysis of which the results are indicated in Table 3

Table 1. List of freshwater gastropods species collected in the field from samples A, B, and C of Fig. 4 within the Bisha mining area, Eritrea

Gastropods species	Sample A	Sample B	Sample C
<i>Melanoides tuberculata</i> (O. F. Müller, 1774)	9	7	4
<i>Bulinus globosus</i> (Morelet, 1866)	7	4	6
<i>Indoplanorbis exustus</i> (Deshayes, 1833)	1	1	0
Total number	17	12	10

Table 2. List of freshwater molluscs and their numbers distinguished from calcareous mud sample from A1 (Fig. 4) within the Bisha mining area, Eritrea

Species name	A1
<i>Melanoides tuberculata</i> (O. F. Müller, 1774)	40
<i>Bulinus globosus</i> (Morelet, 1866)	26
<i>Indoplanorbis exustus</i> (Deshayes, 1833)	56
Bivalve indet.	2



Fig. 5. Closeup picture of locality A1 (Fig. 4) showing the orientation of conical shells approximately parallel to the bedding plane. An arrow indicating the *Indoplanorbis exustus*, occurring less in number at the surface of the mud comparing to the other types of species




Fig. 6. Photos of four species of freshwater shell from A1 at SW 6 within Bisha mining area, Eritrea. A; *Melanoides tuberculata* (O. F. Müller, 1774), B; *Bulinus globosus* (Morelet, 1866), C; *Indoplanorbis exustus* (Deshayes, 1833), D; Bivalve indet

Table 3. Result of ¹⁴C age dating for the two samples of freshwater molluscs from A1 (Fig. 4) within the Bisha mining area, Eritrea

Sample ID	Species	Type of material	δ ¹³ C (‰)	Conventional age (yr BP)	Calibrated age (2σcal BP)	Probability
1a	<i>Indoplanorbis exustus</i>	Shell	-6.06	8270 ± 30	9326-9134	95.4%
1b	<i>Bulinus globosus</i>	Shell	-4.27	8130 ± 30	9134-8999	95.4%

Table 4. Geochemical analysis for the calcareous mud (A1, A2, and A3) from A and soil color in Fig. 4 within Bisha mining area, Eritrea. PAAS and UCC compositions [28] are given for comparison for they represent compositions of earth surface material

Sample	Major elements wt%					Trace elements Sr ppm	TS wt%	Remarks Soil color
	CaO	CaCO ₃	P ₂ O ₅	Fe ₂ O ₃	MnO			
SW-6								
A1	10.04	17.92	0.12	7.24	0.14	483	0.09	10YR 6/2
A2	10.04	17.92	0.13	7.10	0.15	497	0.09	10YR 7/2
A3	8.56	15.28	0.13	7.35	0.14	414	0.09	10YR 7/2
PAAS	1.30	2.32	0.16	6.50				
UCC	3.32	5.93	0.13	4.95	0.10	200	275	




Fig. 7. 3D model geography indicating estimated lacustrine paleoenvironment in western Eritrea. Generated using QGIS, 2018

B. *Bulinus globosus* (Morelet, 1866)

Size: 5–15 mm high and 2–6 mm width

Family: Planorbidae

Ecology: Common habitats are small temporary ponds to large rivers, are shallow waters near shores of lakes, ponds, streams and irrigation channels.

Genus: *Bulinus*

Type species: *Bulinus senegalensis* Müller, 1781

Geographic distribution: Tropical and sub-tropical regions of Africa and Asia.

Synonym: *Physa globosus* Morelet, 1866

Description: *Bulinus globosus* (Morelet, 1866) is oval, whitish in color, has a large aperture with two whorls on their body. Shell length is 5-10 mm while the width is 3-7 mm. The sculpture of the shell consists of faintly growth lines. It has a short conic spire and its coiling direction is sinistral (Fig. 6). The columella of the shells is a bit thin and the inner margin is somehow straight. In some of the shells, the apex is a pit and some are very obtuse.

C. *Indoplanorbis exustus* (Deshayes, 1833)

Order: Heterobranchia

Family: Planorbidae

Genus: *Indoplanorbis*

Type species: *Indoplanorbis exustus*

Synonym: *Planorbis exustus* Deshayes, 1834

Size: 4.5-13 mm high and 5–25 mm wide.

Ecology: These species are found in small ponds, pools, tanks, lakes as well as stagnant pools of water in rivers, and reservoirs. It is also seen less commonly in man-made habitats such as paddy fields, ditches, etc. The snail may also occur in semi-permanent pools formed in flooded areas of fields, where it can survive the dry season buried in mud. It has a life span of average 4 months (India Biodiversity Portal).

Geographic distribution: Widely distributed species; Southeast Asia, Middle East, Africa, India and USA.

Description: *Indoplanorbis exustus* shells have a length that ranges 2-10 mm while the width is 1.4 - 10 mm (Fig. 6). These shells have a dextral coiling direction and whitish in color. The upper side of the shell is evenly rounded, while the back side, the apex is deep but as more whorls grow, it widens up evenly. The number of *Indoplanorbis exustus* increases within the mud. This is because they have invasive nature and ecological tolerance [18] in which it can survive during dry season mingling in the mud.

D. Bivalves indet

Class: Bivalvia

Size: 0.52 mm-1.5 cm long.

Ecology: Bivalves inhabit the tropics, as well as temperate and boreal waters. Live buried in sediments on seabed and sediment in freshwater. A number of species can survive and even flourish in extreme conditions.

Geographic distribution: Due to not identified taxonomy, distribution cannot be described.

Description: The Bivalve shell is 5 mm in width, whitish in color, spherical in shape and consists of equal right and left valves (Fig. 6). The shell sculpture has faint traces of the growth line. The Bivalves were only found buried within the mud. Freshwater bivalves can occur in all latitudes and depth as well as can withstand drought conditions [19].

3.4 ¹⁴C Dating

The ¹⁴C dating was carried out using *Indoplanorbis exustus* and *Bulinus globosus* shells. Result of the ¹⁴C age dating is indicated in Table 3. The analysis was done using AMS in GEOSCIENCE LABORATORY, Nagoya Japan.

3.5 Geochemical Analysis

Abundances of selected major elements (Fe₂O₃* (total iron expressed as Fe₂O₃), MnO, CaO and P₂O₅), Sr and TS (total sulfur) of the calcareous mud samples (A1, A2, and A3 in Fig. 5) were determined by X-ray fluorescence analysis (XRF) in the Department of Earth Science, Shimane University, using a Rigaku Co. Ltd. RIX-2000 spectrometer. Analytical results are indicated in Table 4.

4. DISCUSSION

4.1 Significance of Freshwater Molluscs

Shells molluscs are thin and fragile and are rarely preserved since they usually transport by the water current. Thus, shells reported in this paper could be considered as autochthonous (in situ burial) The abundance of *Melanoides tuberculata* at the surface of the calcareous mud was more compared to the other species. After washing with water, *Indoplanorbis exustus* were found to be more abundant (Table 2), and their sizes were less than 7 mm. The other species were found to have relatively lower concentrations in the mud samples. Bivalve shells were found to occur closed in the left and right valves. This strongly suggests that they were buried alive. Shell preservation is

occasional, and shell bed in the fluvial coarse-grained sediments is very rare. The discovery of shell bed suggested the possibility of a wider distribution of lacustrine deposits in western Eritrea. The results of the ^{14}C dating of both *Indoplanorbis exustus* and *Bulinus globosus*, show 9326-9134 cal yr BP and 9134-8999 cal yr BP, respectively (Table 3). These ages signify the mid-Holocene Epoch, and after 5000 yr BP drastic climatic change was inferred [6]. A number of studies have been carried out on the impact of climatic change after the mid-Holocene Epoch in the East and North African countries by examining Holocene freshwater shells. From the previous data compiled by [20], numerous paleoclimatic data show that in parts of tropical Africa precipitation increased in the early to mid-Holocene which produced the well-documented greening of the Sahara and Sahel, and large lakes of the Holocene have now disappeared in the northern Africa, where higher water levels throughout East Africa were estimated. The discovery of these shells and their ^{14}C age confirmed the existence of a lake in western Eritrea during the mid-Holocene epoch.

Fossilized freshwater molluscs were widespread in tropical and sub-tropical Quaternary deposits [21]. The presence of *Melanoides tuberculata* is important as they were reported to have occurred in the Quaternary deposits throughout Africa and Asia, in both fresh and highly evaporated lakes [22].

Lacustrine deposits are possibly the most powerful basis for integrated insight into the past ecosystems of the world [23]. In contrast to colluvial and alluvial deposits, many lake sediments are characterized by a more or less continuous sedimentary record of the past [24]. Lacustrine deposits are been studied by sedimentological, biological, geochemical and geophysical analyses for quantitative reconstructions of processes of the lakes [25]. The depositional history of lakes is highly sensitive to the interaction of humid and arid climate conditions [26]. Fossil-rich sedimentary deposits provide evidence of the geologic record that often hold chronostratigraphic significance [2]. Fossils of freshwater mollusk shells are among the most common macrofossils from Quaternary lacustrine sediments, which are abundant in calcareous sediments under non-acidic waters [27].

Geochemical composition of calcareous mud samples (A1, A2, A3) showed higher carbonate

(CaCO_3) contents with Sr contents (Table 4) compared to PAAS (post Archean Australian Shale) and UCC (upper continental composition) [28], suggesting enrichment of carbonate matter in this bed. Significantly lower TS (total sulfur) contents showed non-marine depositional condition for the shell bed. Fe_2O_3 contents have slightly higher values than these standards (Table 4) indicating oxic sedimentary conditions. However, MnO and P_2O_5 contents are similar to those of the standard values suggesting a common composition of sediments.

4.2 Holocene Climatic Change and Palaeo-Geography

The mid-Holocene epoch is known as a time when the average temperatures reached the maximum level [5,7,29]. Proxies for rainfall suggest that there was a shift from a predominantly wet episode, known as the African Humid Period (AHP) to dry conditions of the post ca. 5.5 ka [5,7,30,31] in the parts of the northern, eastern and greater Horn of Africa. This African Humid Period came to an abrupt end around 5.5 kyr cal BP [32]. Although the AHP was variable in temporal and spatial distribution, plant and animal communities reorganized into new ecological niches [33,34,35]. The abrupt AHP termination yielded a non-linear, positive biogeophysical feedback between vegetation and precipitation in the Sahara region. The topography of the western part of Eritrea is characterized by landforms of less than 300m above sea level (Fig. 7). There are several small tributary rivers that flow towards the western and southwestern parts of the country, from the highland areas. Accumulated water could form a marsh, which might become a lacustrine environment on the topographic depressions [36]. The outlined depression in Fig. 7 is suggestive of past lake-basin margins estimated to be several tens of km wide. However, this superposition involves inflated assumption. The topographic representations in Fig. 7 could be suggestive of the reconstruction of the western Eritrean mid-Holocene geography.

5. CONCLUSION

This is the first description of the fossilized freshwater molluscus from western Eritrea of which is significant to disclose paleoenvironment and paleoclimate of northeast Africa. The ^{14}C age dating of selected mulluscus confirmed the 8000 yrs BP for the fossil bearing strata composed of calcareous mud. Sedimentary

sequence of this calcareous mud and overlying massive sandy silt to alternating beds of sand and mud demonstrated the lacustrine depositional environment. These lines of fossil evidence and sedimentological feature are considerable information for the paleogeography in western Eritrea during the mid-Holocene. One of the reported mollusc herein, *Melanonids tuberculata* O. F. Müller (1774), a common freshwater species in northern Africa and southern Asia is suggestive of a widespread biota in the huge lake environment in northern Africa. Together with previously supposed lacustrine environment in northern Africa, present report of sediments and fossils enhanced the reconstruction of paleoclimate of warm and humid mode called African Humid Period (AHP). Supposed huge lakes in northern Africa of the mid-Holocene time, however, disappeared due to the climatic changes to dry and hot conditions caused by northeastward expansions of the Sahara Desert.

ACKNOWLEDGEMENT

We thank Japan International Cooperation Agency (JICA) for the scholarship opportunity and Department of Energy and Mines, Eritrea for their permission on exporting the samples and Director General of Department of Mines, Alem Kibreab for his support and kind help.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mandahl-Barth G. Key to the identification of East and Central African freshwater snails of medical and veterinary importance. Bull. Org. mond. Santé, Bull. Wld Hlth Org. 1962;27:135-50.
2. Kidwell SM, Fursich FT, Aigner, T. Conceptual framework for the analysis and classification of fossil concentrations. Palaios. 1986;1(3):228–38. DOI: 10.2307/3514687
3. Cohen AS. Paleolimnology. History and Evolution of Lake Systems. Oxford University Press, New York. 2003;500.
4. Verschuren D, Damsté JSS, Moernaut J, Kristen I, Blaauw M, Fagot M, Haug GH. Half-processional dynamics of monsoon rainfall near the East African Equator. Nature 2009;462(7273):637-41. DOI: 10.1038/nature08520
5. Tierney JE, Lewis SC, Cook BI, LeGrande AN, Schmidt GA. Model, proxy and isotopic perspectives on the East African Humid Period. Earth and Planetary Science Letters. 2011;307(1-2):103-12. DOI: 10.1016/j.epsl.2011.04.038
6. Berke MA, Johnson TC, Werne JP, Grice K, Schouten S, Damsté JS. Molecular records of climate variability and vegetation response since the Late Pleistocene in the Lake Victoria basin, East Africa. Quaternary Science Reviews. 2012a;55:59-74.
7. Berke MA, Johnson TC, Werne JP, Schouten S, Damsté JS. A mid-Holocene thermal maximum at the end of the African Humid Period. Earth and Planetary Science Letters. 2012b;(351–352):95-104.
8. Ghebru B, Araia W, Ogbazghi W, Gebreselassie M, Thomas TS. In East African agriculture and climate change: A comprehensive analysis. Chapter 5. International Food Policy Research Institute (IFPRI). 2013;121-48. Washington DC. Available: <http://ebrary.ifpri.org/cdm/ref/collection/p15738coll2/id/127929>
9. Mineral Potential of the State of Eritrea. Asmara Mining Conference 2014. Ministry of Energy and Mines. Eritrea Mining Journal. The State of Eritrea. 6-11.
10. Barrie CT, Nielsen FW, Aussant CH. The Bisha volcanic-associated massive sulfide deposit, Western Nakfa Terrane, Eritrea. Economic Geology. 2007;102(4):717–38. DOI: 10.2113/gsecongeo.102.4.717
11. Teklay M, Kröner A, Mezger K. Geochemistry, geochronology and isotope geology of Nakfa intrusive rocks, northern Eritrea: Products of a tectonically thickened Neoproterozoic arc crust. Journal of African Earth Sciences. 2001;33(2):283-301. DOI: 10.1016/s0899-5362(01)80064-6
12. Drury SA, Filho CR. Neoproterozoic terrane assemblages in Eritrea: Review and prospects. Journal of African Earth Sciences. 1998;27(3-4):331-48. DOI: 10.1016/s0899-5362(98)00066-9
13. Teklay M. Petrology, Geochemistry and geochronology of neoproterozoic magmatic arc rocks from Eritrea: Implications for crustal evolution in the southern Nubian shield. Department of Mines, Eritrea. 1997; Memoir 1:125.

14. Ogasawara M. Trace element analysis of rock samples by X-ray fluorescence spectrometry, using Rh anode tube. *Bulletin of the Geological Survey of Japan*. 1987;(38):57–68. Japanese with English Abstract.
15. Imai N, Terashima S, Itoh S, Ando A. Compilation of analytical data on nine GSJ geochemical reference samples, "Sedimentary rock series". *Geostandards Newsletter*. 1996;20:165-216.
16. Species summary for *Melanoides tuberculata*". *AnimalBase*, last modified 27 March 2011. (Accessed 19 April 2011) Available:<http://www.animalbase.uni-goettingen.de/zooweb/servlet/AnimalBase/home/species?id=2182>
17. Songtham W, Ugai H, Imsamut S, Maranate S, Tansathien W, Meesook A, Saengsrichan W. Middle Miocene Molluscan Assemblages in Mae Moh Basin, Lampang Provenance, North Thailand. *ScienceAsia*. 2005;(31):183-191. DOI:10.2306/scienceasia1513-1874.2005.31.183
18. Liu L, Mondal M MH, Idris MA, Lokman HS, Rajapakse PRV J, Satrija F, Diaz JL, Upatham ES, Attwood SW. The Phylogeography of *Indoplanorbis exustus* (Gastropoda: Planorbidae) in Asia. *Parasites & Vectors*. 2010;3(1):57. DOI: 10.1186/1756-3305-3-57
19. Morton B. Bivalve. *Encyclopaedia Britannica*; 2018. Available:<https://www.britannica.com/animal/bivalve>
20. Fritz SC. The climate of the Holocene and its landscape and biotic impacts. *Tellus B: Chemical and Physical Meteorology*. 2013;65(1):20602. DOI: 10.3402/tellusb.v65i0.20602
21. Williamson PG. Palaeontological documentation of speciation in Cenozoic mollusks from Turkana Basin. *Nature*. 1981;293(5832):437-43. DOI: 10.1038/293437a0
22. Leng MJ, Lamb AL, Lamb HF, Telford RJ. Palaeoclimatic implications of isotopic data from modern and early Holocene Shells of the freshwater snail *Melanoides tuberculata*, from lakes in the Ethiopian Rift valley. *Journal of Paleolimnology*. 1999;21: 97-106.
23. Oldfield F. Lakes and their drainage basins as units of sediment-based ecological study. *Progress in Physical Geography, Earth and Environment*. 1977;1(3):460-504. DOI: 10.1177/030913337700100303
24. Zolitschka B, Behre KE, Schneider J. Human and climatic impact on the environment as derived from colluvial, fluvial and lacustrine archives—examples from the Bronze Age to the Migration period, Germany. *Quaternary Science Reviews*. 2003;22(1):81-100. DOI: 10.1016/s0277-3791(02)00182-8
25. Zolitschka B, Brauer A, Negendank JFW, Stockjansen H, Lang A. Annually dated late Weichselian continental paleoclimate record from the Eifel, Germany. *Geology*. 2000;28:783-86.
26. Mooij WM, Hu'lsman S, Domis LNDS, Nolet BA, Bodelier PLE, Boers PCM, Pires LMD, Gons HJ, Ibelings BW, Noordhuis R, Portielje R, Wolfstein K, Lammens EHRR. The impact of climate change on lakes in the Netherlands: A review. *Aquatic Ecology*. 2005;39(4):381–400. DOI: 10.1007/s10452-005-9008-0
27. Pukacz A, PeBechaty M, Frankowski M, Kowalski A, Zwijacz-KozsaBka K. Seasonality of Water Chemistry, Carbonate Production, and Biometric Features of Two Species of Chara in a Shallow Clear Water Lake. *The Scientific World Journal Volume*. 2014;1-11. DOI: 10.1155/2014/167631
28. Condie KC. Chemical composition and evolution of the upper continental crust: Contrasting results from surface samples and shales. *Chemical Geology*. 1993; 104(1-4):1-37. DOI: 10.1016/0009-2541(93)90140-e
29. Damsté JSS, Verschuren D, Ossebaar J, Blokker J, Houten RV, van der Meer MT, Plessen B, Schouten S. A 25,000-year record of climate-induced changes in lowland vegetation of eastern equatorial Africa revealed by the stable carbon-isotopic composition of fossil plant leaf waxes. *Earth and Planetary Science Letters*. 2011;302(1-2):236-246.
30. Garcin Y, Junginger A, Melnick D, Olago DO, Strecker MR, Trauth MH. Late Pleistocene–Holocene rise and collapse of Lake Suguta, northern Kenya Rift. *Quaternary Science Reviews*. 2009;28(9-10):911-25.

31. Foerster V, Junginger A, Langkamp O, Gebru T, Asrat A, Umer M, Lamb HF, Wennrich V, Rethemeyer J, Nowaczyk N, Trauth MH, Schaebitz F. Climatic change recorded in the sediments of the Chew Bahir basin, southern Ethiopia, during the last 45,000 years. *Quaternary International*. 2012;274:25-37. DOI: 10.1016/j.quaint.2012.06.028
32. DeMenocal P, Ortiz J, Guilderson T, Adkins J, Sarnthein M, Baker L, Yarusinsky M. Abrupt onset and termination of the African humid period: rapid climate responses to gradual insolation forcing. *Quaternary Science Reviews*. 2000;19(1-5):347-361. DOI: 10.1016/S0277-3791(99)00081-5
33. Gasse F. Hydrological changes in the African tropics since the last glacial maximum. *Quaternary Science Reviews*. 2000;19(1-5):189–211. DOI: 10.1016/S0277-3791(99)00061-X
34. Hély C, Lézine AM, APD contributors. Holocene changes in African vegetation: The tradeoff between climate and water availability. *Climate of the past*. 2014; 10(2):681–86. DOI: 10.5194/cp-10-681-2014.
35. DeMenocal PB. Paleoclimate: End of the African humid period. *Nature Geoscience*. 2015;8(2):86–87. DOI: 10.1038/ngeo2355.
36. Gregory J, Evans DJ, Gebauer S, Howard TG, Hunt DM, Olivero AM, editors. A revised and expanded edition of Carol Reschke's ecological communities of New York State. Second Edition *Ecological Communities of New York State*; 2002.

© 2019 Berhe and Ishiga; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/49443>*