



Response of Integrated Nutrient Management on Irrigated Wheat

S. K. Solanki ^{a*}, H. K. Patel ^a, C. H. Raval ^a and H. K. Patel ^a

^a *Department of Agronomy, B. A. College of Agriculture, Agricultural University, Anand-388110 (Gujarat), India.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i2231353

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/87253>

Original Research Article

Received 15 May 2022
Accepted 21 July 2022
Published 30 July 2022

ABSTRACT

Wheat is one of the most staple foods of the Nepalese diet which is grown in the winter season in most of the area of the country. The study was carried out during *winter* of the year 2019-2020 at College Agronomy Farm, B.A. College of Agriculture, Anand Agricultural University, Anand (Gujarat) to study the response of Integrated Nutrient Management in irrigated wheat under sandy loam soil. Experimental soil is low in organic carbon (0.39 %) and medium in available phosphorus (37.22 kg/ha) and potassium (247 kg/ha) with slightly alkaline (pH 7.85) in reaction. Application of 75% RDN + 25% N through neem cake + *Jeevamrut* (500 lit/ha) treatment reported significantly higher growth attributes and yield attributes like plant height at 30 (25.76 cm), 60 (71.84 cm) DAS and at harvest (119.64 cm), length of spike (10.35 cm), number of effective (305) and total tillers (351), no of grains per spike (49.62). Wheat grain (6389 kg/ha) and straw yield (8950 kg/ha) were significantly higher in 75% RDN + 25% N through neem cake + *Jeevamrut* (500 lit/ha). Crude protein content (13.75 %), soil organic carbon (0.48%) and available phosphorus (53.52 kg/ha) was higher by application of 75% RDN + 25% N through neem cake + *Jeevamrut* (500 lit/ha). Soil microbial population after second and third irrigation was found to be significant by application of 50% RDN + 25% N through neem cake + NP consortium + *Jeevamrut* (500 lit/ha), while after first irrigation response of treatment on soil microbial population, soil pH and EC were found non-significant ($p=0.05$).

*Corresponding author;

Keywords: *Jeevamrut; beejamrut; irrigated wheat; grain yield; neem cake and economics.*

1. INTRODUCTION

Wheat is a key staple food that provides around 20 percent of protein and calories consumed worldwide. Demand for wheat is projected to continue to grow over the coming decades, particularly in the developing world to feed an increasing population and with wheat being a preferred food, continuing to account for a substantial share of human energy needs in 2050.

Integrated nutrient management (INM) is a vital process, and it is the perfect combination of mineral fertilizer, organic manure, compost or bio-fertilizers, and the amalgamation of micronutrients [1]. Integrated nutrient management is practice where all the sources of nutrients namely organic, inorganic and bio-fertilizer as well as liquid organic manures can be combined and applied to soil so that crop growth is enhanced, and we can get good yield with quality product. Liquid organic manure, *jeevamrut* has the potential to play the role of promoting growth and providing resistance in the plant system. Descriptions of this holy combination could be traced to Vedas, the divine scripts of Indian wisdom. Application of *jeevamrut* in agriculture is a good option to supplement nutrient requirement of crops as it is easy to prepare, cost effective, easily available, more reliable and eco-friendlier.

Jeevamrut and *beejamrut* is a liquid organic manure popularly used as means of organic farming as well as integrated nutrient management system. It is excellent source of natural carbon, biomass, Nitrogen, Phosphorus, Potassium and lot of other forms of manure, compost and vermicompost. *Jeevamrut* established to be significant because it maintains the fertility of soil and helps in enhancement of growth and development of plants. *Jeevamrut* also comes in one of the low-cost formulations which are responsible for the enhancement of soil with indigenous micro-organisms required for better mineralization of soil and its helps in enhancement of growth of plant [2,3,4]. Present era is on integrated nutrient system and we have to observed all possible side with its benefit for successful crop production. View of these objects present studies on integrated nutrient management was taken and it included combined used organic manures, liquid fermented manures, chemical fertilizers and

micro organized and its response on irrigated wheat.

2. MATERIALS AND MATHODS

A field experiment was conducted at College Agronomy Farm, B. A. College of Agriculture, Anand Agricultural University, Anand (Gujarat) during *winter* of the year 2019-20 to study the effect of integrated nutrient management in irrigated wheat. The soil of the experimental field was loamy sand in texture, medium in organic carbon (0.39 %) and medium in available phosphorus (37.22 kg/ha) and available potash (247 kg/ha) with slightly alkaline (pH 7.85) in reaction. Wheat variety Gujarat Wheat 451 (GW 451) was selected for experiment. The experiment was arranged in randomized block design with three replications, consisting of ten treatments **T₁**: 100% RDN through chemical fertilizer, **T₂**: 75% RDN + 25% N through neem cake, **T₃**: 75% RDN + 25% N through neem cake + NP consortium, **T₄**: 75% RDN + 25% N through neem cake + *Jeevamrut* (500 lit/ha), **T₅**: 50% RDN+ 50% N through neem cake, **T₆**: 50% RDN + 50% N through neem cake + NP consortium, **T₇**: 50% RDN + 50% N through neem cake + *Jeevamrut* (500 lit/ha), **T₈**: 50% RDN + 25% N through neem cake + NP consortium + *Jeevamrut* (500 lit/ha), **T₉**: 50% RDN + NP consortium + *Jeevamrut* (500 lit/ha), **T₁₀**: NP consortium + *Jeevamrut* (500 lit/ha). Recommended nitrogen (120 kg N/ha) was given in form of Urea at 50% as a basal and remaining 50% nitrogen in two splits at CRI (Crown root initiation) and tillering stage while recommended dose of phosphorus (60 kg P₂O₅/ha) in the form of Single Super Phosphate was applied as a basal. Application of bio NP consortium with seed treatment (5 ml/kg) and soil application (1 lit/ha) with first three irrigation as per treatments. Application of *jeevamrut* (500 lit/ha) at first three irrigation as per treatment. After sowing crops and proper germination in all plots, selected and tagged five random plant in net plot area for further growth and yield observations. The protein content of grain was assessed by multiplying nitrogen content of the seed (%) with the conversion factor of 6.25. Soil sample were collected after five days of irrigation applied for first three irrigation plot wise separately and stored in polythene bags and kept at 4°C till processed. Microbial count done by serial dilution technique (Dhingra and Sinclair, 1993) and spreaded on nutrient agar (NA) plat and

incubated at 30-35° C for 2-3 days in incubator. Colonies were counted by using colonies counter meter and total microbial count were taken by following formula;

Final count (cfu/g) = (Number of well isolated colonies x Dilution factor / Aliquot taken)

The collected data for various parameters were statistically analysed using Fishers analysis of variance (ANOVA) technique and the treatments were compared at 5% levels of significance [5]. All the observation, growth and yield parameters were taken as per standard method.

3. RESULTS AND DISCUSSION

Response of different integrated nutrient management treatments on plant population in meter row length was found non-significant ($p=0.05$) at 20 DAS, it might be due to uniform germination in all treatments, at initial stage response of nutrient was found non-significant, so uniformity was observed. Application of 75 % RDN + 25 % N through neem cake + *Jeevamrut* (500 lit/ha) reported significantly ($p=0.05$) higher plant at 30, 60 DAS and at harvest (25.76,71.84 and 119.64 cm, respectively). Significantly ($p=0.05$) higher plant might be due to organic manure could be attributed to slow release of nitrogen and increased availability of micro and macro nutrients available in neem cake to produce new meristematic tissues. Besides, NPK and micronutrients are also available in neem cake which plays a vital role in various metabolic activities of plants and catalytic role in activating several enzymes [6, 7 and 8]. The liquid organic manures contain microbial population and plant growth promoting substances that help in improving plant growth, metabolic activities of plants.

Application of 75% RDN + 25% N through neem cake + *Jeevamrut* (500 lit/ha) reported significantly higher length of spike (10.35 cm), effective and total tillers per meter row length (305 and 351, respectively), number of spikes per plant at harvest (10.40) and number of grains per spikes (49.62) but it did not differ significantly ($p=0.05$) with treatment T_1 (100% RDN through chemical fertilizer). Response on growth attributing characters might be due to combined effect of organic manure, biofertilizers and chemical fertilizers in balanced proportion played a very important role in decomposition and easy release of different nutrients and their uptake by the crop which result into higher dry matter

accumulation and its translocation in different plant parts of growth and yield parameters, which resulted into higher yield. *Jeevamrut* content major and micronutrient which are rapidly available and easily absorbed and helps in faster growth and development of plant component. Similar line of results was also reported by [9, 10, and 11] (Fig.02).

Data presented in Fig.03 revealed that significantly ($p=0.05$) higher grain yield (6389 kg/ha) and straw yield (8950 kg/ha) was observed significantly higher in T_4 (75% RDN + 25% N through neem cake + *Jeevamrut* (500 lit/ha) treatment. Treatment T_4 (75% RDN + 25% N through neem cake + *Jeevamrut* (500 lit/ha) reported 6.16% higher grain yield over 100 % RDF treatment (T_1). Significantly higher grain and straw yield by application of organic and inorganic nutrients might be due to supply of balanced amount of essential nutrients to plants in suitable proportion and forms. It is well established fact that the incorporation of organic manure not only acts as a store house of major and micronutrients but is also influence the physical, chemical and biological properties of soil [12]. Organic manure contains almost all the essential plant nutrients, its incorporation in soil in conjunction with inorganic fertilizers promotes rapid vegetative growth and tillering, thereby, increasing the sink size in terms of ear and grain growth. The application liquid manure supplied the required metabolites to the source and further greater accumulation of assimilates in the sink [11, 13] reported that *jeevamrut* can release the nutrients in a more synchronized manner as per the need of the crops and *jeevamrut* enhances microbial activity in soil and helps in mobilization of nutrient in soil. Other reason might be due to combined effect of organic manure (FYM), biofertilizers and chemical fertilizers with sulphur in balanced proportion played a very vital role in decomposition and easy release of different nutrients and their uptake by the crop which led to higher dry matter accumulation and its translocation in different plant parts of growth and yield parameters, which in turn resulted into higher yield [9].

Protein content (13.75 %) was found to be significantly ($p=0.05$) higher in treatment T_4 (75% RDN + 25% N through neem cake + *Jeevamrut* (500 lit/ha) than rest of integrated nutrient management treatments except treatment T_1 (100% RDN through chemical fertilizer). The response of treatment on soil pH and EC did not differ significantly ($p=0.05$) by effect of various




Fig. 1. Effect of treatments on growth parameters of wheat




Fig. 2. Effect of treatment on yield parameters of wheat




Fig. 3. Effect of treatment on grain and straw yield of wheat




Fig. 4. Effect of treatment on economics of wheat

Table 1. Effect of treatment on protein content, soil microbial count and soil properties after harvest of crop

Treatment	Grain protein content (%)	Soil microbial count ($\times 10^6$ cfu/g)			pH	EC	OC (%)	P ₂ O ₅
		After 1 st irrigation	After 2 nd irrigation	After 3 rd irrigation				
T ₁	13.23	81.52	111.52	119.85	8.05	0.25	0.47	52.51
T ₂	11.76	85.67	125.67	150.67	8.05	0.24	0.46	44.68
T ₃	11.86	103.53	143.53	168.53	8.02	0.26	0.46	47.07
T ₄	13.75	92.89	132.89	157.89	8.02	0.25	0.48	53.52
T ₅	9.34	89.79	129.79	154.79	7.98	0.22	0.39	39.91
T ₆	11.19	103.87	143.87	168.87	7.95	0.23	0.41	40.22
T ₇	11.39	101.39	141.39	166.39	7.99	0.24	0.45	43.12
T ₈	11.60	108.44	148.44	173.44	8.00	0.25	0.45	43.62
T ₉	9.29	107.88	147.88	172.88	8.03	0.27	0.39	38.99
T ₁₀	9.12	95.2	138.54	163.54	7.99	0.27	0.38	37.51
S. Em. \pm	0.57	6.23	6.78	7.29	0.26	0.01	0.02	1.86
CD(P=0.05)	1.69	NS	20.13	21.66	NS	NS	0.05	5.54
CV %	8.76	11.12	8.61	7.91	5.61	8.85	6.51	7.32

treatments. Application of 75% RDN + 25% N through neem cake + *Jeevamrut* (500 lit/ha (T₄)) reported higher organic carbon (0.48 %) and available P₂O₅ (53.52 kg/ha). Soil microbial population was found non-significant (p=0.05) at first irrigation, but it was significantly (p=0.05) higher during second (148.44 $\times 10^6$ cfu/g) and third irrigation (173.44 $\times 10^6$ cfu/g) under treatment 50% RDN + 25% N through neem cake + NP consortium + *Jeevamrut* (500 lit/ha). Increasing microbial population might be due to availability of abundant organic matter and effective microbial activities because of sufficient

supply of feeding material for microorganism in the form of humus. Similar finding reported by Pawar et al. [14].

4. ECONOMICS

Based on present market prices of wheat seed and straw as well as different variable and non-variable inputs. The data on cost of cultivation, gross and net return as well as BCR were calculated for different treatments are presented in Fig.04. Application of treatment T₁ (100% RDN through chemical fertilizer) recorded higher net

realization (Rs. 95907/ha) and BCR value (4.6) followed by application of treatment T₄ [75% RDN + 25% N through neem cake + Jeevamrut (500 lit/ha)] recorded net realization (Rs.82677/ha) and BCR value (2.76). However, lower net realization and BCR value recorded under application of treatment T₅ (50% RDN+ 50% N through neem cake).

4. CONCLUSION

Based on the above results, it can be concluded that the grain yield and economics of wheat can be improved by using different integrated nutrient management treatment. Experiment results also revealed that wheat morphological and physiological parameters improved by application of organic and inorganic with liquid manure.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Sibananda D, Manoj S, Langyan S, Geeta S, Rakesh P, Avadhesh S et al. Integrated nutrient management reduced the nutrient losses and increased crop yield in irrigated wheat. Arch Agron Soil Sci. 2022;1-12.
- Singh AS, Lal EP. Impact of organic liquid formulation, jeevamrutha on photosynthetic pigments of *Ocimum basilicum* L. (sweet basil) under NaCl induced salinity stress. Plant Arch. 2019;19(2):1997-2001.
- Devakumar N, Rao GGE, Shubha S, Imrankhan N, Gowda SB. Activities of organic farming research centre. Shimoga: Navile. Bangalore: University Agricultural Science; 2008.
- Sreenivasa MN, Naik N, Bhat SN. Beejamrutha: A source for beneficial bacteria. Karnataka J Agric Sci. 2010;22(5):1038-40.
- Steel RG, Torrie JH. Principles and procedures of statistics. New Delhi: McGraw-Hill Book Company-110 001; 1982.
- Shivakumar BC, Girish AC, Gowda B, Kumar GCV, Gowda APM, Thimmegowda MN. Influence of Pongamia, mahua and neem cakes on finger millet productivity and soil fertility. J Appl Nat Sci. 2011;3(2):274-6.
- Verma SB, Singh D, Chauhan RM. Effect of neem (*Azadirachta indica* L.), mustard (*Brassica juncea*) de-oiled seed cake and bio-fertilizer on the growth and yield of wheat (*Triticum aestivum* L.). J Pharmacogn Phytochem. 2018;7(5):2416-27.
- Neelam RK, Nanwal RK, Kumar P. Effect of organic and inorganic sources of nutrients on productivity and profitability of mungbean-wheat cropping system. Legu Rese - An Inter Jour. 2015;38(4):509-12.
- Desai HA, Dodia IN, Desai CK, Patel MD, Patel HK. Integrated nutrient management in wheat (*Triticum aestivum* L.). Trends Biosci. 2015;8(2):472-5.
- Safiullah K, Durani A, Durrani H, Ansari MA. Effect of solid and liquid organic manures on growth, yield and economics of sweet corn (*Zea mays* L.) under south Gujarat condition. International Journal of Pure and Applied Bioscience. 2018;6(2):567-74.
- Dalvi SM, Ghodpage RM, Balpande SS, Badole WP, Patil SS, Sirsat DD. Effect of integrated plant nutrient system on phosphorus solubilisation and productivity of wheat in vertisols. J Pharmacogn Phytochem. 2020;9(6):765-9.
- Yadav VK, Oury F, Suda N, Liu ZW, Gao XB, Confavreux C et al. A serotonin-dependent mechanism explains the leptin regulation of bone mass, appetite, and energy expenditure. Cell. 2009 Sep 4;138(5):976-89.
- Patel JS, Patel GJ, Patel KM, Patel BM. Effect of drip irrigation and micronutrient mixture on growth, yield and quality of sweet corn. Curr Adv Agric Sci. 2013; 5(1):124-5.
- Pawar VR, Tambe AD, Patil SP, Suryawanshi SU. Effect of different organic inputs on yield, economics and microbial count of sweet corn (*Zea mays* Var. Saccharata). Ecol Environ Conserv. 2013; 19(3):865-8.

© 2022 Solanki et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/87253>