

# International STD Research & Reviews 3(1): 1-7, 2015; Article no.ISRR.2015.001 ISSN: 2347-5196



## SCIENCEDOMAIN international

www.sciencedomain.org

# Assessing Health Education Techniques in Enhancing the Knowledge of HIV/AIDS among Adolescents

Ashok Pandey<sup>1\*</sup>, S. Umashankar<sup>1</sup>, Hai I. Dai<sup>1</sup>, Chetraj Pandit<sup>1</sup>, Min Kunwar<sup>1</sup>, Madhu Pandey<sup>2</sup> and Natisara Rai<sup>3</sup>

<sup>1</sup>Padmashree School of Public Health, Rajiv Gandhi University of Health Sciences (RGUHS), Nagarbhavi Circle, Bangalore-560072, Karnataka, India. <sup>2</sup>Pashupati Multiple Campus, Chabahil, Kathmandu, Nepal. <sup>3</sup>Shakti Milan Samaj (SMS), Dhumbahari, Kathmandu, Nepal.

#### Authors' contributions

This work was carried out in collaboration between all authors. All authors read and approved the final manuscript.

#### Article Information

DOI: 10.9734/ISRR/2015/10961

Editor(s)

(1) Constantinos Petrovas, Immunology Laboratory, Vaccine Research Center, NIAID/NIH, Bethesda, USA.
(2) Gabriella GD D'ettorre, Department of Public Health and Infectious Diseases, University of Rome "Sapienza" and Azienda
Policlinico Umberto I, Italy.

(3) Ma Luo, National Microbiology Lab., Public Health Agency of Canada, Canada and Department of Medical Microbiology, University of Manitoba, Canada.

i, Cariaui Poviowor

(1) Anonymous, Nigeria.

(2) Anonymous. Melaka Manipal Medical College. Malaysia.

Complete Peer review History: http://www.sciencedomain.org/review-history.php?iid=646&id=27&aid=6244

Original Research Article

Received 18<sup>th</sup> April 2014 Accepted 19<sup>th</sup> July 2014 Published 27<sup>th</sup> September 2014

#### **ABSTRACT**

**Introduction:** Adolescent refers to individuals between the ages of 10-19 years. In Nepal, Adolescent comprises more than 22% of population. Educations are important as a 'social vaccine', and it can serve as a powerful preventive tool.

**Methods:** The study was conducted on three secondary school of in Hansapur Village Development Committees, Arghakhanchi district. The sampling design used for the study was stratified random sampling. A sample size of 300 adolescent students was taken.

**Results:** Mean pre-intervention knowledge scores were 57.36±17.44 for pre intervention groups. After health education by the five methods in the five subgroups, the pooled mean knowledge score was enhanced to 81.80±16.47. It was highly significant (p<0.001). The overall increase in Post-

\*Corresponding author: E-mail: pandeyg7@gmail.com;

intervention mean score in the intervention group (From Pre- intervention to Post-intervention) was 20.32 percent. The corresponding pre-intervention mean scores were 10.98±5.04, 12.06+6.01, 12.35±5.68, 10.98±4.99, and 10.98±4.99 respectively. Immediately after the educational activities the mean knowledge scores (Post-intervention score) enhanced to 13.13±4.96, 14.93±5.96, 16.16±6.25, 19.36±5.50, 18.20±7.16 in the book, lecture, poster pamphlets, video and participatory lecture groups respectively. It was highly significant for all the five intervention subgroups.

**Conclusion:** Video and the participatory lecture are the most effective health education techniques for effective delivery of HIV/AIDS. It is suggested that programme implementers might chose the suitable methods required for their individual programmes.

Keywords: Adolescents; effectiveness; health education program; knowledge.

#### 1. INTRODUCTION

Generally, the term "adolescent" refers to individuals between the ages of 10-19 years. Adolescent is a period of physical, psychological and social maturing from childhood to adulthood. Because of the physical, social and psychological transition adolescents face many different health risks and on the other hand. Such curiosity and interest in learning offers great opportunities for improving adolescent health and development [1].

In Nepal, Adolescent comprises more than 22% of population. The Median age at marriage is 16.8 years for girls and three years later in men. Median age at first sexual intercourse for women is nearly identical to their median age at first marriage however men tend to initiate sex about one year before marriage. In case of HIV, 13% of all HIV cases are adolescents aged 14-19 years and 70% of them are females. There were 20% of estimated 25,000 client sex workers are under the age of 16 years. In case of drug use young people 16-19 years constituted 22.5% of the total population. In Nepal. the topography. environmental degradation, poverty economic migration are all linked and they combine with other factors to increase vulnerability to HIV. For reduce vulnerability educational methods were effective in enhancing the knowledge of the students [2].

Three educational methods namely lecture, participatory; pamphlets were effective in enhancing the knowledge of the students [3]. Adolescents have excellent resources for delivering effective education: skilled teachers; an interactive educational process that occurs over time; a variety of learning opportunities; materials and methods; and the ability to involve parents in their children's learning [4]. Among various methods of disease prevention health education finds a very significant role". Different

types of health education methods create awareness in the community as well as organization, Educations is important as a 'social vaccine', and it can serve as a powerful preventive tool [5].

Education is one of the pillars of development and providing universal access to primary education by 2015 is one of the Millennium Development Goals. HIV infection as a good basic education ranks among the most effective and cost-effective means of preventing HIV [6]. The assessing health education techniques in health education research are seldom seen since the concept is relatively new to the health sector. The National Health Policy of Nepal intends to target school children and adolescents for promoting healthy behaviors among the general population (MoHP, 1991). Hence, the present study expresses the various processes of these health education interventions and to understand the education for better methods of Health education techniques.

### 1.1 Objective of the Study

To assess the various health education methods in enhancing the knowledge of HIV/AIDS among adolescents.

#### 2. MATERIALS AND METHODS

The study was conducted on three secondary school of in Hansapur Village Development Committees, Arghakhanchi district. The school was selected by convenience sampling methods. The sampling design used for the study was stratified random sampling (Fig. 1).

Self-administered questionnaire was used for collecting data (Fig. 2).

This sample size was found to be statistically adequate. The content of the health education

was kept similar for the five intervention groups and was delivered by one person only. The data was analyzed using the Statistical Package for Social Sciences (SPSS) Version 16 (SPSS for Windows 16.0, SPSS Inc., Chicago, IL) and Microsoft office package 2010. The data has been analyzed using mean and percentages, and paired t test was used to test the improvement in knowledge. Validity and reliability has been maintained by pre-test and a necessary modification has been made. Consultation has been done with the supervisor/guide/subject experts. The data were collected with full voluntary verbal and written consent and not forcefully. It had been taken care of full confidentiality. They were assured that all the information belonging to the respondent will be kept confidential and data will be utilized only for the purpose of this study. Informed consent was obtained from all the respondents before data collection.

# 2.1 Research Hypotheses

- "There is no difference in the overall knowledge score of adolescent students about HIV/AIDS between pre intervention and post intervention".
- "There is no difference in the knowledge of adolescent students about HIV/AIDS between pre intervention and post intervention of different techniques of health education methods".




Fig. 1. Schematic design of sampling technique




Fig. 2. Flow chart of tools and techniques

#### 3. RESULTS

It is observed from the present study that 61 % of male and 39 % of female were taken for this survey. The study reveals that majority of respondents 92.3 % were unmarried and 7.7 were married. Similarly, it is found that 35.7 % of the respondents were Chhetri, 23.3 % were Janajati, 20 % were Dalit and 20 % were Brahmins. The study reveals that almost all of the respondents 99% were Hindu and 1% was Christians (Table 1).

Mean pre-intervention knowledge mean scores were 57.36  $\pm 17.44$  for pre intervention groups. After health education by the five methods in the five subgroups, the pooled mean knowledge score was enhanced to 81.80  $\pm 16.47$  (Post-intervention score). It was highly significant (P = 0.001). The overall increase in post-intervention mean score in the intervention group (From Pre-intervention to Post-intervention) was 20.32 percent (Table 2).

The total number of students (both the intervention) in the book, lecture, poster pamphlets, video and participatory lecture groups were 60 each. The corresponding preintervention mean scores were 10.98±5.04,

12.06+6.01, 12.35±5.68, 10.98±4.99, and 10.98±4.99 respectively. Immediately after the educational activities the mean knowledge scores (Post-intervention score) enhanced to 13.13±4.96, 14.93±5.96, 16.16±6.25, 19.36±5.50, 18.20±7.16 in the book, lecture, poster pamphlets, video and participatory lecture groups each. This immediate increase in knowledge scores from pre to post-intervention. It was highly significant for all the five intervention subgroups (Table 3).

'Sustainable increase' in knowledge was calculated as the percentage of increased knowledge from pre-intervention to post-intervention. The differences in the differences in the pre-intervention and post intervention mean knowledge scores between the five groups were statistically significant. It was calculated to be 21.5, 28.7, 38.1, 84.16, and 72.4 percent respectively for lecture, poster pamphlets, video and participatory lecture (Table 4).

Table 1. Socio-demographic characteristics of respondents

Socio-demographic characteristics			
Male	183	61.0	
Female	117	39.0	
Total	300	100.0	
Married	23	7.7	
Unmarried	277	92.3	
Total	300	100.0	
Brahmins	60	20.0	
Chhetri	107	35.7	
Janajati	70	23.3	
Dalit	63	21.0	
Total	300	100.0	
Hindu	297	99.0	
Christian	3	1.0	
Total	300	100.0	
	Male Female Total Married Unmarried Total Brahmins Chhetri Janajati Dalit Total Hindu Christian	Male       183         Female       117         Total       300         Married       23         Unmarried       277         Total       300         Brahmins       60         Chhetri       107         Janajati       70         Dalit       63         Total       300         Hindu       297         Christian       3	

Table 2. Mean knowledge scores of students in pre intervention and post intervention groups

Intervention	N(300)	Mean + Std. deviation	Percentage increase in knowledge score*	t- value	P-value
Pre intervention	300	57.36+17.44	100.8	25.02	0.000
Post intervention	300	81.80+16.47	121.12	37.99	0.000

<sup>\*</sup> Percentage of increase score retained from pre-intervention to post- intervention

Table 3. Comparison of mean knowledge scores in five interventions group

Intervention group	ention group Pre interven		Post intervention		Percentage increase in	
	N	Mean+Std. deviation	N	Mean+Std. deviation	─ knowledge*	
Book	60	10.98+5.04	60	13.13+4.96	21.5	
Lecture	60	12.06+6.01	60	14.93+5.96	28.7	
Poster/pamphlets	60	12.35+5.68	60	16.16+6.25	38.1	
Video	60	10.98+4.99	60	19.36+5.50	84.16	
Participatory	60	10.98+4.99	60	18.20+7.16	72.4	

<sup>\*</sup> Percentage of increase score retained from pre-intervention to post- intervention

Table 4. Test of significance within the same intervention subgroups

Groups	Pre-intervention vs post-intervention			
	Mean+Std. deviation	T	P value	
Book	-2.150+5.89	-2.824	0.006	
Lecture	-2.86+6.62	-3.350	0.001	
Poster/ pamphlets	-3.81+6.79	-4.353	0.000	
Video	-8.38+7.56	-8.582	0.000	
Participatory lecture	-7.21+8.25	-6.770	0.000	

#### 4. DISCUSSION

The findings of the study showed that education intervention played an important role in increasing knowledge about HIV/AIDS which can be supported by the study "Evaluation of a School Based HIV/AIDS Educational Intervention in Ukraine" [7].

In present study reveals that the 13% increase in knowledge of the post intervention after intervention of different health education technique. Therefore, we can say that educational methods have been able to enhance the knowledge after intervention by 13 percent. We observe that there is significant difference between pre-intervention and post-intervention knowledge scores.

HIV/AIDS, which might be the reasons behind relatively limited knowledge of the respondents about HIV/AIDS before education intervention and after intervention there was a significant increase in respondents' knowledge. The study findings indicated that there was significant increase in the knowledge in all the aspects of HIV/AIDS after education intervention. All of the respondents indicated that there is a need of awareness program about HIV/AIDS. This finding is supported by participatory method of teaching there is active participation of the participants and they learn more and are able to retain the gained knowledge [3].

In this present study participatory lecture and video method is more effective as compared to read, lecture and poster/ pamphlet methods because in the participatory method of teaching there is active participation of the participants and they learn more and are able to retain the gained knowledge. And in video methods more sense organ are concentrate to seeing, listening. Post-intervention score for book, lecture and poster/pamphlet group was significantly less than the other two groups. This could be because of the fact that despite our motivation it is quite likely that many of the students either did not read or partially read. And in listen students are used to the daily lecture. Although knowledge was better than practice, both were not satisfactory. So, the adolescent should be educated about the process and significance of HIV/AIDS [8].

The increased gain in knowledge was sustained to a very high degree. It was approximately

similar for book, lecture, poster/pamphlets (21.5, 28.7 and 38.1 percent's respectively) and much higher for video and participatory lecture group (84.16 and 72.4 percent). The reason for this high sustainability in the pamphlet group could be because of the reason that once the students were sensitized to a particular issue they were more sensitive to related information through various mass media and other IEC activities going on in the area.

#### 5. CONCLUSION

From the above discussion it can be concluded that education played an important role in increasing knowledge of the respondents about HIV/AIDS. In view of the above results and observation, it is concluded that all the five educational methods were effective in enhancing the knowledge of the students. But Video and the participatory lecture are the most effective health education techniques for effective delivery of HIV/AIDS. It is suggested that programme implementers might chose the suitable methods required for their individual programmes. However, it is not ever suggested that these are the only methods available.

#### **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

#### **REFERENCES**

- Rao Rs. Effectiveness of reproductive health education among rural adolescent girls: a school based intervention study in Udupi taluk, Karnataka Indian J Med Sci. 2008;62(11):439-443.
- Population Division MoHPGoN Kathmandu Nepal, Nepal Demographic and Health Survey 2011, Population Division: New ERA Kathmandu, Nepal, ICF International Calverton, Maryland, U.S.A; 2012.
- Singh SP, et al. Effect of some health educational techniques in disseminating the knowledge about HIV/ AIDS among adolescent student. Indian Journal of Community Medicine. 2005;30(1)1-3.
- Kayode AD, Ogu MA. School location and secondary school students' awareness of Human Immuno-Deficiency Virus/Acquired Immune Deficiency Syndrome (HIV/AIDS) In Imo State of Nigeria.

- Journal of Educational and Social Resear ch. 2011;1(2):39-47.
- 5. Srivastava AN, Mahmood SY, Mahmood PA, et al. Adolescence awareness: A better tool to combat HIV/AIDS. Journal of National Community Medicine. 2011;2(1):86-90.
- 6. Luncheon Remarks on Women and AIDS at Interaction Forum by Geeta Rao Gupta, PhD. International Center for Research on Women; 2005.
- 7. Kyrychenko P, Kohler C, Sathiakumar N. Evaluation of a school- based HIV/AIDS educational intervention in Ukraine. J Adolesc Health. 2006;39(6):900-7. Epub 2006 Aug 30.
- 8. Pandey A. Challenges experienced by adolescent girls while menstruation in Kathmandu, Valley. A qualitative study IOSR-JNHS. 2014;3(2):41-45.

© 2015 Pandey et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

#### Peer-review history:

The peer review history for this paper can be accessed here: http://www.sciencedomain.org/review-history.php?iid=646&id=27&aid=6244